




Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

15. 04. 2004

Best Available Copy



Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03005181.7

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



Anmeldung Nr.:
Application no.: 03005181.7
Demande no:

Anmeldetag:
Date of filing: 07.03.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

BASF Plant Science GmbH

67056 Ludwigshafen
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Enhanced amylose production in plants

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT SE SI SK TR LI

Enhanced amylose production in plants

Description

5 The present invention relates to new amylose biosynthesis enhancing proteins, nucleic acids encoding a starch biosynthesis enhancing protein, a method for producing amylose with high efficiency by culturing genetically modified plants with an increased amylose biosynthesis compared to the wild type or to the genetically modified plants themselves as well as the use of these transgenic plants over-expressing at least one
10 of the amylose biosynthesis enhancing proteins for the production of amylose.

Starch is the major storage carbohydrate of plants and is mainly accumulated in seeds and tubers, which are then the reproductive tissues of plants that form those types of organs. Starch is also accumulated on a diurnal basis where starch is built up in green
15 tissue from photosynthetic products and then metabolised for energy during the dark period. The storage starch is assembled into semi crystalline granules. Amylopectin and amylose are the two constituent molecules of starch. Amylopectin is a branched molecule consisting of linear α -1,4 glucan chains linked by α -1,6 bonds. Amylose consists essentially of the linear α -1,4 glucan chains.
20

Starch is utilised for many applications within the technical industry as well as the food industry. Main crops used by starch processors are maize and potato. For potatoes specific varieties are utilised for starch production that have been bred for high starch contents. This means that the starch content and yield is an important economic driver
25 for the starch processing industry. A greater part of produced dry starch is used for paper production. The specifications and requirements for the starch component varies from application to application and starch is many times chemically modified in order to provide desired properties to an application. Another way to achieve starch of different qualities is to take advantage of mutations in the starch biosynthesis and more recently
30 by genetic modification of pathways leading to starch. The first main modifications have been to separate the production of the two starch components amylopectin and amylose into different varieties. Waxy or "amylose free" varieties contain solely amylopectin type starch while there are also high amylose genotypes such as "amylose extender" in maize.
35

Amylose starch has several potential industrial uses as a film former or for expanded products. High amylose starch can be achieved in potatoes and other starch containing plants by inhibition of starch branching enzymes. This leads then to the concomitant reduction or elimination of amylopectin branching and thereby an increased amylose
40 fraction.

US 5,856,467 describes the genetically engineered modification of potato for suppressing formation of amylopectin-type starch. The document describes an antisense construct for inhibiting, to a varying extent, the expression of the gene coding for formation of starch branching enzyme (SBE gene) in potato, said antisense construct comprising a tuber specific promoter, transcription start and the first exon of the SBE gene, inserted in the antisense direction.

US 6,169,226 relates to an amino acid sequence of a second starch branching enzyme (SBE II) of potato and a fragment thereof as well as to the corresponding isolated DNA sequences. It describes the production of transgenic potatoes and the use of these transgenic potatoes for the production of amylose-type starch.

WO 97/20040 and WO 98/20145 describe methods of altering the amylopectine/amylose starch content of plant cells by introducing into the plant cells nucleic acid sequences operably linked in sense or antisense orientation to a suitable promoter which homologous genes encodes polypeptides having SBE I or SBE II activity.

A side effect of the amylose overproduction is a decreased total starch content in the potatoes. This decrease becomes more pronounced as the amylose fraction is increased.

Basic enzymes for the production of amylopectin and amylose are starch synthases that build the linear α -1,4 glucan chains and branching enzymes breaking the α -1,4 glucan chain and reattaching them by α -1,6 bonds. Several other enzymes are likely to affect starch structure and composition, such as debranching enzymes, but initially most focus has been towards affecting the expression of starch synthases and starch branching enzymes. This has led to an extensive dissection of what enzymes are important for what features of starch synthesis. However it has never been convincingly shown how the synthesis of starch in plants whether amylose or amylopectin is initiated.

Suggestions on the initiation of starch biosynthesis have been the subject of several scientific papers since it has been difficult to attribute a primer independent function to starch synthases under other than artificial in vitro conditions. By primer independent function implies the formation of new α -1,4 glucan chains with ADP-glucose as the sole starting point and building block. One proposed pathway has been that the presence of maltooligosaccharides act as primers for the addition of further glucose units by starch synthases although it has been debated on whether concentrations are sufficient to provide the basis for starch synthesis and also how these maltooligosaccharides would be formed in the plastids.

Starch is in plants synthesised as an energy storage molecule. Much is known about the enzymes participating in the starch biosynthesis although, the initiation of the starch molecule. In mammals and yeast an energy storage molecule very similar to starch is synthesised, glycogen. The enzymatic steps for synthesis of the respective molecules are analogous. In glycogen biosynthesis the initiation of the molecule is known and synthesised by the enzyme glycogenin. Glycogenin is a self-glucosylating enzyme polymerising a linear chain of approximately 8 glucose molecules on itself. The primer of about 8 glucose residues is necessary for the enzymes catalysing the continuation of glucose incorporation to the glycogen molecule to function.

Cheng et al., 1995, Mol. and Cell. Biol. 6632-6640 compare the two yeast proteins with rabbit muscle glycogenin.

Roach et al., 1997, Progress in Nucleic Acid Research and Molecular Biology Vol 57, describe self glycosylating initiator proteins and their roll in glycogen biosynthesis.

Mu et al., 1997, Journal of Biological Chemistry 272 (44), 27589-27597 compare mammalian with yeast and C. elegans glycogenins.

Factors important for starch quantity have been investigated and many initiatives have been taken, especially in potato, to increase starch formation and content by over-expression or inhibition of various enzyme activities in areas of increased substrate supply, increased biosynthesis activity or shutting down substrate diverting pathways but so far this has led only to limited success with no commercial applications and only some scientific publications.

Regierer, B. et al., Starch content and yield increase as a result of altering adenylate pools in transgenic plants. Nat Biotechnol. 20(12):1256-60, (2002).

Sweetlove, LJ et al., Starch synthesis in transgenic potato tubers with increased 3-phosphoglyceric acid content as a consequence of increased 6-phosphofructokinase activity. Planta 213(3):478-82 (2001).

Veramendi, J et al., Antisense repression of hexokinase 1 leads to an overaccumulation of starch in leaves of transgenic potato plants but not to significant changes in tuber carbohydrate metabolism. Plant Physiol. 121(1):123-34 (1999).

Geigenberger, P et al., Overexpression of pyrophosphatase leads to increased sucrose degradation and starch synthesis, increased activities of enzymes for sucrose-

starch interconversions, and increased levels of nucleotides in growing potato tubers. *Planta*. 205(3):428-37(1998).

- 5 Sweetlove, LJ et al., Starch metabolism in tubers of transgenic potato (*Solanum tuberosum*) with increased ADPglucose pyrophosphorylase. *Biochem J.* 320 (Pt 2):493-8 (1996).

10 In other research a biochemical function superficially similar to the one initiating glycogen production in animals was investigated. A class of genes have then been isolated from several plants and was given the name amylogenin (WO94/04693; Sing, D. et al; β -Glucosylarginin: a new glucose-protein bond in a self-glucosylating protein from sweet corn, *FEBS Letters* 376:61-64, (1995) in the belief that it was the plant equivalent of glycogenin which acts as a self-glucosylating enzyme and provide primers for starch biosynthesis in plants. These genes have no resemblance from a structural point
15 of view to the genes coding for glycogenin and have later been determined not to have a function in starch biosynthesis but rather might be of importance for cell wall formation, see Bocca, S.N et al., Molecular cloning and characterization of the enzyme UDP-glucose: protein transglucosylase from potato. *Plant Physiology and Biochemistry* 37(11):809-819(1999).

20 WO 98/50553 describes nucleic acid fragments encoding a plant glycogenin or a water stress protein. WO 98/50553 also relates to the construction of chimeric genes encoding all or a portion of a plant glycogenin in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of a plant glycogenin in a transformed host cell.
25

Thus although many enzymes and pathways have been investigated in plants, the question on how starch formation is initiated and what determines the starch content is still unresolved.

30 Amylose is a commercially important starch product with many uses but unfortunately an increase in amylose content in transgenic potato plants is associated with a significant decrease in starch content, see figure 1.

35 Analyses of transgenic high amylose potato lines show that there is an excess of soluble sugars in these lines, see figure 2. This indicates that the starch biosynthesis in these transgenic lines is not efficient enough for incorporation of available sugars.

40 Amylose starch consists of very few reducing ends compared to native starch. Therefore it is commercially important to identify genes that further enhance the amylose

biosynthesis and that are capable to incorporate the excess of glucose residues available and to compensate the decrease in starch content in plants that produces amylose in high amounts.

- 5 The invention aims at enhancing the yield of amylose biosynthesis by the over-expression of genes which enhance amylose biosynthesis in transformed plants.

The invention describes genes coding for proteins which enhance amylose production.

- 10 The present invention describes the nucleic acids SEQ ID NO 1 and 3 from potato coding for enzymes enhancing the de novo amylose biosynthesis.

- 15 Example 1 describes that the nucleic acid sequences SEQ ID NO 1 or 3 can complement a missing glycogenin function in yeast cells containing knock-out mutations for the self-glycosylating proteins Glg1p and Glg2p.

- 20 Gene constructs were made for gene-inhibition and over-expression of the two genes SEQ ID NO 1 or 3 in potato. Transgenic lines with the over-expressed or inhibited enzyme activity were analysed with regard to the genes influence on amylose content.

Both genes were inserted in sense and antisense direction downstream of a plant promoter element, resulting in the transformation binary vectors pHS1, pHS2, pHS3 and pHS4, see figures 3-7.

- 25 The antisense constructs were transformed into potato plant varieties Prevalent and Producent and the sense constructs were transformed to the potato varieties Desirée and the transgenic plant AM99-2003 according to the transformation method as described in example 2. The transgenic plant AM99-2003 was produced as described in example 3.

- 30 Prevalent and Producent are starch varieties having a starch content of approximately 20 %. Desirée is a potato variety having a starch content of approximately 16% and AM99-2003 is a transgenic high amylose line having a starch and thereby amylose content of approximately 13%.

- 35 The putative genes were isolated from a tuber specific cDNA library of *Solanum tuberosum* (variety Prevalent). The library was made from a lambdaZAP directional kit (Stratagene).

Both cDNAs isolated were full-length clones of the individual genes and named StGH1 and StGH2, for nucleic acid sequences see SEQ ID NO 1 and SEQ ID NO 3.

pHS1

5

A 1300bp PCR fragment from the StGH1 gene was constructed in antisense direction driven by the gbss promoter. The PCR fragment was cut out from its cloning vector pCR4-TOPO (Invitrogen) with EcoRI (blunted) and XbaI. The fragment was ligated to the pGPTV-kan (Becker, D. et al., Plant Molecular Biology 20:1195-1197(1992) based binary vector pHo3.1 between a gbss promoter (WO 92/11376) and a nos terminator at the Sall (blunted) and XbaI sites. The binary vector also includes nptII as selection marker driven by the nos promoter (Herrera, L. et al., 1983). The construct was named pHS1, for details see figure 3a and 4.

15 pHS2

A 2300bp full-length cDNA clone of StGH2 was cut out from the cloning vector pBluescript (Stratagene) with XbaI and XhoI. The gene was ligated in antisense direction between the gbss promoter and nos terminator to the binary vector pHo3.1 at XbaI and Sall. As can be seen under pHS1 the vector has nptII as selection system. The vector was named pHS2, for details see figure 3b and 5.

pHS3

25 A full-length StGH1 cDNA, (1780bp) was cut out from the host vector pBluescript with EcoRI (blunted) and BglII and ligated to the BamHI and SmaI sites of pUCgbssprom (3886bp), containing pUC19 with the gbss promoter and the nos terminator. The plasmid was named pUCGH1.

30 A fragment with the gbss promoter, the StGH1 gene and the nos terminator was moved from pUCGH1 with EcoRI (blunted) and HindIII (2980bp) and ligated to PstI (blunted) and HindIII opened pSUN1 (WO 02/00900). The plasmid was named pSUNGH1.

35 A 3600bp fragment containing the AHAS resistance gene from Arabidopsis thaliana (Sathasivan, K. et al., Plant Physiology 97(1991), 1044-1050) with nos promoter, see Herrera-Estrella, L. et al., Nature 303:209-213(1983) and OCS terminator (Wesley, S.V. et al., Plant J. 27(6):581-590(2001) was ligated to pSUNGH1 (9000bp) at the SmaI site. The vector was given the name pHS3, for details see figure 3c and 6.

pHS4.

The gbss promoter and nos terminator was ligated to pBR322 with EcoRI and HindIII. Between the promoter and terminator an EcoRI-HincII full-length gene pStGH2 was cloned at the XbaI site. The 3366bp promoter-gene-terminator complex was cut using EcoRI (partial digestion) and EcoRV, and ligated to pSUN1 at EcoRI-EcoRV and named pSUNGH2. An XbaI fragment with AHAS gene (*Arabidopsis thaliana*), nos promoter and OCS terminator was ligated to pSUNGH2 opened with XbaI (partial digestion). The AHAS gene is used as selection marker. The construct was named pHS4, for details see figure 3d and 7.

Example 2 describes the general method for the transformation of different potato plant varieties producing native starch or high amylose type starch with pHS1, pHS2, pHS3 or pHS4.

The StGH1 and StGH2 genes were down-regulated in the potato plant varieties Prevalent and Producent by transformation with the genes in antisense direction in relation to a plant regulatory element as described in example 4 and 6. Down-regulation of the two genes resulted in a decrease in gene expression in transgenic lines, compared to their mother varieties in the order of 50-95%, see example 7 and table 3. Transgenic lines transformed with pHS1 and pHS2 with confirmed decrease in gene-expression have a decrease in dry matter of 7 to 11% compared to their mother varieties, see example 8 and table 5.

The StGH1 and StGH2 genes were over-expressed in potato driven by the tuber specific promoter gbss, as described in example 5. A mutated AHAS gene was used as selection marker yielding tolerance to the Imazamox herbicides. Two potato varieties were transformed, Desiree and AM99-2003 a transgenic high amylose line with a 40% decrease in starch content compared to its mother variety. The transformed lines over-expressing StGH1 and StGH2 were selected as described in example 6. The gene expression levels were analysed with real-time PCR, see example 7 and table 3. The over-expression of the genes StGH1 and StGH2 resulted in an 2 to 10 times increase in gene compared to their mother variety. Furthermore the lines over-expressing StGH1 and StGH2 showed an increase in dry matter of up to 36 % as described in example 8 and table 5.

The over-expression of StGH1 and StGH2 in transgenic potato plants producing amylose type starch resulted in an increased dry matter content, which means an increased amylose content as no amylopectine is produced, see example 8, 9 and 11.

The amylose biosynthesis enhancing protein according to the invention comprises the amino acid sequence SEQ ID NO 2 or 4 or a protein which comprises a sequence derived from SEQ ID NO 2 or 4, which is at least 50%, preferably at least 60%, more preferably at least 70%, more preferably at least 80%, still more preferably at least 90%, most preferably at least 95%, identical at the amino acid level to the sequence SEQ ID NO 2 or 4 and has the property of an amylose biosynthesis enhancing protein. This amylose biosynthesis enhancing protein may also be prepared by artificial variations starting from the SEQ ID NO 2 or 4, for example by substitution, insertion or deletion of amino acids.

Such a protein could also be used to increase the production of starch or amylopectin in non-transgenic or transgenic plants.

The term "substitution" in the specification means the replacement of one or more amino acids by one or more amino acids. Preference is given to carrying out "conservative" replacements in which the amino acids replaced has a property similar to that of the original amino acid, for example replacement of Glu by Asp, Gln by Asn, Val by Ile, Leu by Ile, Ser by Thr.

"Deletion" is the replacement of an amino acid or amino acids by a direct bond. Preferred positions for deletions are the polypeptide termini and the junctions between the individual protein domains.

"Insertions" are insertions of amino acids into the polypeptide chain, with a direct bond formally being replaced by one or more amino acids.

"Identity" between two proteins means the identity of the amino acids over the in each case entire length of the protein, in particular the identity which is calculated by comparison with the aid of the Vector NTI Suite 7.1 Software of the company Informax (USA) using the Clustal W method (Thompson, JD et al., Nucleic Acid Research, 22 (22):4673-4680, 1994)

with the parameters set as follows:

Multiple alignment parameter:

Gap opening penalty	15
Gap extension penalty	6.66
Gap separation penalty range	8
Gap separation penalty	on

% identity for alignment delay	40
Residue specific gaps	on
Hydrophilic residue gap	off
Transition weighing	0

5

Pairwise alignment parameter:

FAST algorithm	off
K-tuple size	2

10

Gap penalty	5
Window size	4
Number of best diagonals	4

15 Accordingly, a protein which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4 means a protein which, when comparing its sequence with the sequence SEQ ID NO 2 or 4, is at least 50% identical, in particular according to the above program algorithm using the above set of parameters.

20 Further natural examples of genes coding for an amylose enhancing protein according to the invention can readily be found, for example, in various organisms, in particular in plants, whose genomic sequence is known by comparing the identity of the amino acid sequences or of the corresponding back-translated nucleic acid sequences from data-bases with the sequence of SEQ ID NO 2 or 4, in particular according to the above .

25 program algorithm using the above set of parameters.

In the completed genome sequence of *Arabidopsis thaliana*, five putative coding sequences can be deduced by searching for exon/intron boundaries and comparing with back translated sequences of SEQ ID NO 2 or 4.

30

The following nucleic acid sequences of *Arabidopsis thaliana* SEQ ID NO 5, SEQ ID NO 7, SEQ ID NO 9, SEQ ID NO 11 and SEQ ID NO 13 could be used to carry out the invention and are coding for the amylose biosynthesis enhancing proteins SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12 and SEQ ID NO 14.

35

Furthermore the following nucleic acid sequences or ESTs can be used in order to identify and clone genes coding for an amylose biosynthesis enhancing protein from plant organisms:

Tomato ESTs from GenBank: AW216407, BE450055, BF097262, BE450557, BF097173

5 Wheat ESTs from GenBank: BJ292476, BJ278875, BJ283925, BE442966, CA666180, BQ483228

Maize EST from GenBank: BG319971

10 Rice ESTs from GenBank: AL606633, CA752890, BI813265

Natural examples of amylose biosynthesis enhancing proteins and the corresponding genes can furthermore readily be found in various organisms, in particular plants, whose genomic sequence is unknown by hybridization techniques in a manner known per se, for example starting from the nucleic acid sequences SEQ ID NO 1 or SEQ ID
15 NO 3 or any of the SEQ ID NO 5, 7, 9, 11 or 13 or any of the EST sequences described above.

The hybridization may be carried out under moderate (low stringency) or, preferably, under stringent (high stringency) conditions.

20 Such hybridization conditions are described, inter alia, in Sambrook, J., Fritsch, E.F., Maniatis, T., in: Molecular Cloning (A Laboratory Manual), 2nd edition, Cold Spring Harbor Laboratory Press, 1989, pages 9.31-9.57 or in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6.

25 By way of example, the conditions during the washing step may be selected from the range of conditions which is limited by those with low stringency (with 2X SSC at 50°C) and those with high stringency (with 0.2X SSC at 50°C, preferably at 65°C) (20X SSC: 0.3 M sodium citrate, 3 M sodium chloride, pH 7.0).

30 In addition, the temperature may be raised during the washing step from moderate conditions at room temperature, 22°C, to stringent conditions at 65°C.

35 Both parameters, salt concentration and temperature, may be varied simultaneously and it is also possible to keep one of the two parameters constant and to vary only the other one. It is also possible to use denaturing agents such as, for example, formamide or SDS during hybridization. In the presence of 50% formamide, the hybridization is preferably carried out at 42°C.

40 Some exemplary conditions for hybridization and washing step are listed below:

(1) hybridization conditions with, for example

5 (i) 4X SSC at 65°C, or

(ii) 6X SSC at 45°C, or

(iii) 6X SSC at 68°C, 100 mg/ml denatured fish sperm DNA, or

10 (iv) 6X SSC, 0.5% SDS, 100 mg/ml denatured fragmented salmon sperm DNA at 68°C, or

(v) 6X SSC, 0.5% SDS, 100 mg/ml denatured fragmented salmon sperm DNA, 50% formamide at 42°C, or

15

(vi) 50% formamide, 4X SSC at 42°C, or

(vii) 50% (vol/vol) formamide, 0.1% bovine serum albumin, 0.1% Ficoll, 0.1% polyvinylpyrrolidone, 50 mM sodium phosphate buffer pH 6.5, 750 mM NaCl, 75 mM sodium citrate at 42°C, or

20

(viii) 2X or 4X SSC at 50°C (moderate conditions), or

(ix) 30 to 40% formamide, 2X or 4X SSC at 42°C (moderate conditions).

25

(2) Washing steps of 10 minutes each with, for example

(i) 0.015 M NaCl/0.0015 M sodium citrate/0.1% SDS at 50°C, or

30 (ii) 0.1X SSC at 65°C, or

(iii) 0.1X SSC, 0.5% SDS at 68°C, or

(iv) 0.1X SSC, 0.5% SDS, 50% formamide at 42°C, or

35

(v) 0.2X SSC, 0.1% SDS at 42°C, or

(vi) 2X SSC at 65°C (moderate conditions).

Preferred proteins with amylose biosynthesis enhancing activity are proteins from plants, cyanobacteria, mosses or algae, particular preferred from plants. A particular preferred protein comprises the amino acid sequence SEQ ID NO 2 or 4.

- 5 If, for example, the protein is to be expressed in a plant, it is frequently advantageous to use the codon usage of said plant for backtranslation and resynthesis of the gene according to codon usage of said plant.

- 10 The invention further relates to nucleic acids encoding an amylose biosynthesis enhancing protein according to the invention. All of the nucleic acids mentioned in the specification may be, for example, a RNA sequence, DNA sequence or cDNA sequence.

- 15 Suitable nucleic acid sequences can be obtained, for example, by back-translating the polypeptide sequence according to the genetic code. For this, preference is given to using those codons which are used frequently according to the organism-specific codon usage. The codon usage can be readily determined on the basis of computer analyses of other known genes of the organisms in question.

- 20 All of the above-mentioned genes coding for an amylose biosynthesis enhancing protein can furthermore be prepared in a manner known per se from the nucleotide building blocks by chemical synthesis, for example by fragment condensation of individual overlapping complementary nucleic acid building blocks of the double helix. The chemical synthesis of oligonucleotides may be carried out, for example, in a known
25 manner according to the phosphoramidite method (Voet, Voet, 2nd edition, Wiley Press New York, pages 896-897). The annealing of synthetic oligonucleotides and filling-in of gaps with the aid of the Klenow fragment of DNA polymerase and ligation reactions and also general cloning methods are described in Sambrook et al. (1989), Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press.

- 30 Genes coding for this function may be integrated in the plant chromosomes and upon expression utilize a transit peptide to localise to plastids which is the organelle where amylose biosynthesis takes place or be integrated directly into the plastid genome and thereby surpass the need for the localisation signal. The genes may be expressed
35 constitutively or organ specific. For organ specific expression, promoters with tuber specific expression is preferable in potatoes while in cereals as maize or wheat a endosperm specific expression would be preferred to achieve a high degree of expression in organs where storage starch is accumulated. When transformed to the plastid genome then specific regulatory elements suitable for that organelle apply.

The genes of this invention may be used in combination with other genes that can be situated on the same gene construct or transferred and combined by co-transformation or super transformation. Genes and traits that are of interest to combine with the genes of the instant invention are agronomic or input trait such as herbicide tolerance, disease and pest resistance or stress tolerance but could also be output traits such as starch structure modification or yield. Genes and traits used in combination with the genes described in the invention could be for adding a function that is not present in the modified plant species or over-expressing a function that is already present or inhibiting a function by the use of antisense, RNAi or antibodies.

The invention may be used to increase the amylose content in potato tubers but would in its context not be limited to potatoes but would be applicable to other starch producing and storing plants such as e.g. corn, cassava, wheat, barley, oat and rice.

The described invention is particularly suited for eliminating a lower starch content associated with increased amylose content in different plants where the number of α -1,4-glucan chain non-reducing ends is greatly reduced due to the reduction or elimination of α -1,6 branch formation. Amylopectin is an extremely efficient structure, as is glycogen, for polysaccharide production since it is very branched and thus contains as many points accessible for starch synthesis as there are non-reducing ends. Starch that is mainly composed of amylose, contains much fewer branches and thus the biosynthetic capacity is reduced. In order to enhance starch biosynthesis when there is no amylopectin production, expression of genes as described in the present invention, could for example form new primers that can replace amylopectin as a source for starch biosynthesis capacity and thereby reduce or eliminate the lost capacity for starch synthesis. To further illustrate the situation the degree of branching in ordinary potato starch is approximately 3.1% while in high amylose starch it is 0.3-1.0% depending on amylose content this decrease of branching and starch content is further associated with an increase in glucose and fructose content.

The increased amylose content and thereby solids content is also advantageous for the processing properties in various applications such as for french fries, potato crisps and other potato based products. In addition to an increased solid content, the inserted genes SEQ ID NO 1 or 3 of the present invention result in the transformation of excess sugars into α -1,4-glucan chains and thereby reducing browning of fried potato products, Maillard reaction, in which amino acids react with free sugars.

Furthermore

- (i) any gene of plant origin with the described activity can be used for increasing amylose content and solids.

- (ii) the genes can be controlled by any regulating promoter element functional in plant.
- (iii) any starch producing crop of any variety can be transformed with the described genes.
- 5 (iiii) any plant transformation method can be used.
- (iiii) any binary vector can be used for the insertion of the described genes.
- (iiii) the described genes can be combined with any other desired transgenically inserted traits.
- 10 The invention further relates to a method for producing amylose by culturing plants which have, compared to a wild type or a genetically modified plant producing already amylose type starch, an increased amylose biosynthesis activity, said proteins comprising the amino acid sequence SEQ ID NO 2 or 4 or a sequence which is derived from one of these sequences by substitution, insertion or deletion of amino acids and which
- 15 is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4.
- Increased amylose biosynthesis activity compared to the wild type or transgenic line means that the amount of amylose formed is increased by the amylose biosynthesis enhancing protein in comparison with the wild type or transgenic line.
- 20 This increase in amylose biosynthesis activity is preferably at least 5%, further preferably at least 10%, further preferably at least 20%, further preferably at least 50%, more preferably at least 100%, still more preferably at least 200%, in particular at least 500%, of the protein activity of the wild type or transgenic line.
- 25 A "wild type" means the corresponding genetically unmodified starting plant. This plant is preferably *Solanum tuberosum*.
- 30 Depending on the context, the term "plant" means a wild type starting plant or a genetically modified starting plant.
- "Transgenic plant" or "genetically modified plant" means that the plant contains an additional inserted gene segment that may be foreign or endogenous to the plant species, additional genes or additional gene fragments in sense and/or antisense orientation to a suitable promoter corresponding to the following polypeptides and showing enzymatic activity of a starch branching enzyme I, a starch branching enzyme II and/or the amylose biosynthesis enhancing protein as specified in SEQ ID NO 1 or 3 or polynucleotides having at least 60 % sequence identity thereof.
- 35

"Amylose type starch" means that the amylose content of the starch is increased compared to the amylose content of starch produced by wild type plants especially wild type potato plants.

- 5 The amylose biosynthesis activity may be increased in various ways, for example by eliminating inhibiting regulatory mechanisms at the translation and protein levels or by increasing the gene expression of a nucleic acid encoding a amylose biosynthesis enhancing protein compared to the wild type or transgenic plant, for example by inducing a gene encoding the amylose biosynthesis enhancing protein via activators or by
10 introducing into the plant nucleic acids encoding a amylose biosynthesis enhancing protein.

According to the invention, increasing the gene expression of a nucleic acid encoding a amylose biosynthesis enhancing protein could also mean manipulating the expression
15 of the endogenous amylose biosynthesis enhancing protein intrinsic to the plant, in particular in potato plants. This may be achieved, for example, by modifying the promoter DNA sequence of genes encoding an amylose biosynthesis enhancing protein. Such a modification which leads to a modified or preferably increased rate of expression of at least one endogenous gene encoding an amylose biosynthesis enhancing
20 protein may be carried out by deleting or inserting DNA sequences.

It is also possible to modify expression of one or more endogenous amylose biosynthesis enhancing protein by applying exogenous stimuli. This may be carried out by particular physiological conditions, i.e. by applying foreign substances.
25

Furthermore, it is possible to achieve a modified or increased expression of at least one endogenous gene encoding an amylose biosynthesis enhancing protein by the interaction of a regulatory protein which is modified or is not present in the untransformed plant.
30

In a preferred embodiment, the amylose biosynthesis enhancing protein activity is increased compared to the wild type or transgenic plant by increasing the gene expression of a nucleic acid encoding an amylose biosynthesis enhancing protein, said amylose biosynthesis enhancing protein comprising the amino acid sequence SEQ ID NO
35 2 or 4 or a sequence which is derived from said sequences by substitution, insertion or deletion of amino acids and which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4.

In the case of genomic nucleic acid sequences encoding an amylose biosynthesis enhancing protein from eukaryotic sources, which contain introns, preferably already
40

processed nucleic acid sequences such as the corresponding cDNAs are to be used, if the host organism is unable to or cannot be enabled to express the corresponding amylose biosynthesis enhancing protein.

- 5 In this preferred embodiment, the transgenic plant of the invention thus contains, compared to the wild type or transgenic plant, at least one further gene encoding an amylose biosynthesis enhancing protein. In this preferred embodiment, the genetically modified plant of the invention has accordingly at least one transgenic endogenous or exogenous nucleic acid encoding an amylose biosynthesis enhancing.

10

Suitable and preferred nucleic acids are described above. In a particularly preferred embodiment, a nucleic acid comprising the sequence SEQ ID NO 1 or 3 is introduced into the plant.

- 15 According to the invention, organisms means preferably eukaryotic organisms, such as, for example, yeasts, algae, mosses; fungi or plants, which are capable of producing starch or amylose, either as wild type or enabled by genetic modification. Preferred organisms are photosynthetically active organisms such as, for example, plants which, even as a wild type, are capable of producing starch or amylose type starch.

20

Particularly preferred organisms are potato plants.

- 25 The present invention furthermore relates to the use of proteins comprising the amino acid sequence SEQ ID NO 2 or 4 or a sequence which is derived from this sequence by substitution, insertion or deletion of amino acids and which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4 and having amylose biosynthesis enhancing activity.

- 30 The present invention further relates to the use of nucleic acids SEQ ID NO 1 or 3 or one of the SEQ ID NOs 5, 7, 9, 11 or 13 encoding proteins having an amylose biosynthesis enhancing activity in plants.

- 35 The transgenic organisms, in particular plants, are preferably prepared by transforming the starting organisms, in particular plants, with a nucleic acid construct containing the above-described nucleic acid, encoding an amylose biosynthesis enhancing protein which is functionally linked to one or more regulatory signals ensuring transcription and translation in said organisms.

These nucleic acid constructs in which the coding nucleic acid sequence is functionally linked to one or more regulatory signals ensuring transcription and translation in organisms, in particular in plants, are also referred to as expression cassettes herein below.

- 5 Accordingly, the invention further relates to nucleic acid constructs, in particular to nucleic acid constructs functioning as expression cassette, which comprise a nucleic acid encoding an amylose biosynthesis enhancing protein which is functionally linked to one or more regulatory signals ensuring transcription and translation in organisms, in particular in plants.

10

The regulatory signals preferably comprise one or more promoters ensuring transcription and translation in organisms, in particular in plants.

- 15 The expression cassettes include regulatory signals, i.e. regulatory nucleic acid sequences, which control expression of the coding sequence in the host cell. According to a preferred embodiment, an expression cassette comprises upstream, i.e. at the 5' end of the coding sequence, a promoter and downstream, i.e. at the 3' end, a polyadenylation signal and, where appropriate, further regulatory elements which are operatively linked to the coding sequence for at least one of the above-described genes
- 20 located in between. Operative linkage means the sequential arrangement of promoter, coding sequence, terminator and, where appropriate, further regulatory elements in such a way that each of the regulatory elements can properly carry out its function in the expression of the coding sequence.

- 25 When the organism used is a plant, the nucleic acid constructs and expression cassettes of the invention preferably contain a nucleic acid encoding a plastid transit peptide ensuring localisation in plastids.

- 30 The preferred nucleic acid constructs, expression cassettes and vectors for plants and methods for preparing transgenic plants and also the transgenic plants themselves are described in examples 2 to 6 below.

- 35 The sequences preferred for operative linkage, but not limited thereto, are targeting sequences for ensuring subcellular localisation to plastids such as amyloplasts or chloroplasts but could also mean in the apoplasts, in the vacuole, in the mitochondrion, in the endoplasmic reticulum (ER), in the nucleus, in elaioplasts or in other compartments and translation enhancers such as the tobacco mosaic virus 5'-leader sequence (Gallie et al., Nucl. Acids Res. 15 (1987), 8693-8711).

A suitable promoter of the expression cassette is in principle any promoter which is able to control the expression of foreign genes in plants.

- 5 "Constitutive" promoter means those promoters which ensure expression in numerous, preferably all, tissues over a relatively long period of plant development, preferably during the entire plant development.

- 10 Preference is given to using, in particular, a promoter from plants or a promoter originating from a plant virus. Preference is in particular given to the promoter of the 35S transcript of the CaMV cauliflower mosaic virus (Franck et al. (1980) Cell 21:285-294; Odell et al. (1985) Nature 313:810-812; Shewmaker et al. (1985) Virology 140:281-288; Gardner et al. (1986) Plant Mol Biol 6:221-228) or the 19S CaMV promoter (US 5,352,605; WO 84/02913; Benfey et al. (1989) EMBO J 8:2195-2202).

- 15 Another suitable constitutive promoter is the Rubisco small subunit (SSU) promoter (US 4,962,028), the leguminB promoter (GenBank Acc. No. X03677), the Agrobacterium nopaline synthase promoter, the TR double promoter, the agrobacterium OCS (octopine synthase) promoter, the ubiquitin promoter (Holtorf S et al. (1995) Plant Mol Biol 29:637-649), the ubiquitin 1 promoter (Christensen et al. (1992) Plant Mol Biol 20 18:675-689; Bruce et al. (1989) Proc Natl Acad Sci USA 86:9692-9696), the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (US 5,683,439), the promoters of the vacuolar ATPase subunits or the promoter of a proline-rich wheat protein (WO 91/13991), the Pnit promoter (Y07648.L, Hillebrand et al. (1998), Plant. Mol. Biol. 36, 89-99, Hillebrand et al. (1996), Gene, 170, 197-200) and other promoters of genes 25 whose constitutive expression in plants is known to the skilled worker.

- The expression cassettes may also contain a chemically inducible promoter (review: Gatz et al. (1997) Annu Rev Plant Physiol Plant Mol Biol 48:89-108) which may be used to control expression of the amylose biosynthesis enhancing protein gene in the 30 plants at a particular time. Promoters of this kind, such as, for example, the PRP1 promoter (Ward et al. (1993) Plant Mol Biol 22:361-366), salicylic acid-inducible promoter (WO 95/19443), a benzenesulfonamide-inducible promoter (EP 0 388 186), a tetracycline-inducible promoter (Gatz et al. (1992) Plant J 2:397-404), an abscisic acid-inducible promoter (EP 0 335 528) and an ethanol- or cyclohexanone-inducible promoter 35 (WO 93/21334), may likewise be used.

- Further examples of suitable promoters are fruit ripening-specific promoters such as, for example, the fruit ripening-specific promoter from tomato (WO 94/21794, EP 409 625). Development-dependent promoters partly include the tissue-specific promoters, 40 since individual tissues are naturally formed in a development-dependent manner.

Furthermore, preference is given in particular to those promoters which ensure expression in tissues or parts of the plant, in which, for example, biosynthesis of starch or amylose or of the precursors thereof takes place. Preference is given, for example, to promoters with specificities for leaves, stems, roots, seeds and tubers.

- Seed-specific promoters are, for example, the phaseoline promoter (US 5,504,200; Bustos MM et al. (1989) Plant Cell 1(9):839-53); the promoter of the 2S albumin gene (Joseffson LG et al. (1987) J Biol Chem 262:12196-12201), the legumin promoter (Shirsat A et al. (1989) Mol Gen Genet 215(2): 326-331), the USP (unknown seed protein) promoter (Bäumlein H et al. (1991) Mol Gen Genet 225(3):459-67), the promoter of the napin gene (US 5,608,152; Stalberg K et al. (1996) L Planta 199:515-519), the sucrose-binding protein promoter (WO 00/26388) and the legumin B4 promoter (LeB4; Bäumlein H et al. (1991) Mol Gen Genet 225: 121-128; Baeumlein et al. (1992) Plant Journal 2(2):233-9; Fiedler U et al. (1995) Biotechnology (NY) 13(10):1090f), the Arabidopsis oleosin promoter (WO 98/45461), the Brassica Bce4 promoter (WO 91/13980) and the vicillin promoter (Weschke et al. 1988, Biochem. Physiol. Pflanzen 183, 233-242; Bäumlein H et al. (1991) Mol Gen Genet 225(3):459-67).
- Further suitable seed-specific promoters are those of the genes coding for high molecular weight glutenine (HMWG), gliadin, branching enzyme, ADP glucose pyrophosphatase (AGPase) and starch synthase. Preference is further given to promoters which allow seed-specific expression in monocotyledons such as e.g. corn, barley, wheat, rye, rice, etc. It is also possible to use advantageously the promoter of the *lpt2* or *lpt1* gene (WO 95/15389, WO 95/23230) or the promoters described in WO 99/16890 (promoters of the hordein gene, the glutelin gene, the oryzin gene, the prolamin gene, the gliadin gene, the glutelin gene, the zein gene, the kasirin gene and the secalin gene).
- Examples of tuber-, storage root- or root-specific promoters are the patatin promoter class I (B33), the potato cathepsin D inhibitor promoter and the potato granular bound starch synthase (GBSS) promoter as described in EP-A 0 921 191.
- Examples of leaf-specific promoters are the cytosolic FBPase promoter from potato (WO 97/05900), the rubisco (ribulose-1,5-bisphosphate carboxylate) SSU (small subunit) promoter and the potato ST-LSI promoter (Stockhaus et al. (1989) EMBO J 8:2445-2451).

Further promoters suitable for expression in plants have been described (Rogers et al. (1987) Meth in Enzymol 153:253-277; Schardl et al. (1987) Gene 61:1-11; Berger et al. (1989) Proc Natl Acad Sci USA 86:8402-8406).

5 The site of starch and amylose biosynthesis in potato plants is the amyloplast. Therefore amyloplast-specific targeting and activity of the gene products of the inventivenucleic acids SEQ ID NO 1 or 3 encoding an amylose biosynthesis enhancing protein is desirable.

10 The expression may also take place in a tissue-specific manner in all parts of the plant.

A further preferred embodiment therefore relates to a tuber-specific expression of the nucleic acids SEQ ID NO 1 or 3.

15 In addition, a constitutive expression of the gene encoding an amylose biosynthesis enhancing protein is advantageous. On the other hand, however, an inducible expression of this gene may also be desirable.

20 An expression cassette is preferably prepared by fusing a suitable promoter to an above-described nucleic acid encoding an amylose biosynthesis enhancing protein and, preferably, to a nucleic acid which has been inserted between promoter and nucleic acid sequence and which codes for an amyloplast-specific transit peptide and also to a polyadenylation signal according to familiar recombination and cloning techniques as described, for example, in T. Maniatis, E.F. Fritsch and J. Sambrook, Molecular
25 Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989) and in T.J. Silhavy, M.L. Berman and L.W. Enquist, Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1984) and in Ausubel, F.M. et al., Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley-Interscience (1987).

30 Particular preference is given to inserted nucleic acid sequences which ensure targeting in the amyloplasts.

35 It is also possible to use an expression cassette in which the nucleic acid sequence encodes an amylose biosynthesis enhancing protein fusion protein, one part of the fusion protein being a transit peptide which controls translocation of the polypeptide. Preference is given to amyloplast-specific transit peptides which, after translocation of amylose biosynthesis enhancing protein into the amyloplasts, are enzymatically
40 cleaved off the amylose biosynthesis enhancing protein part.

Particular preference is given to the transit peptide which is derived from the *Nicotiana tabacum* plastid transketolase or from another transit peptide (e.g. the transit peptide of the rubisco small subunit or of ferredoxin NADP oxidoreductase and also of isopentenyl pyrophosphate isomerase-2) or from its functional equivalent.

5

Further examples of a plastid transit peptide are the transit peptide of the plastid isopentenyl pyrophosphate isomerase-2 (IPP-2) from *Arabidopsis thaliana* and the transit peptide of the ribulose biphosphate carboxylase small subunit (rbcS) from pea (Guerineau, F, Woolston, S, Brooks, L, Mullineaux, P (1988) An expression cassette for targeting foreign proteins into the chloroplasts. Nucl. Acids Res. 16: 11380).

10

Plant genes of the invention which encode a plant amylose biosynthesis enhancing protein may already contain the nucleic acid sequence which encodes a plastid transit peptide. In this case, a further transit peptide is not required. For example, the *Solanum tuberosum* sequences of the amylose biosynthesis enhancing protein of the invention SEQ ID NO 1 or 3 contain already a transit peptide sequence.

15

The nucleic acids of the invention may be prepared synthetically or obtained naturally or comprise a mixture of synthetic and natural nucleic acid components and may also be composed of various heterologous gene sections of various organisms.

20

As described above, preference is given to synthetic nucleotide sequences with codons which are preferred by plants. These codons which are preferred by plants may be determined from codons which have the highest frequency in proteins and which are expressed in most of the interesting plant species.

25

When preparing an expression cassette, it is possible to manipulate various DNA fragments in order to obtain a nucleotide sequence which expediently can be read in the correct direction and is provided with a correct reading frame. The DNA fragments may be linked to one another by attaching adaptors or linkers to said fragments.

30

It is furthermore possible to use manipulations which provide appropriate restriction cleavage sites or which remove excess DNA or restriction cleavage sites. In those cases for which insertions, deletions or substitutions such as, for example, transitions and transversions are suitable, in vitro mutagenesis, primer repair, restriction or ligation can be used.

35

Preferred polyadenylation signals are polyadenylation signals functional in plants, exemplified by those which correspond essentially to T-DNA polyadenylation signals from *Agrobacterium tumefaciens*, in particular of the T-DNA gene 3 (octopine synthase).

40

or OCS terminator, the complete sequence of the Ti plasmid pTiACH5 (Gielen et al., EMBO J. 3, 835 -846(1984) or functional equivalents.

- 5 The invention further relates to the use of the nucleic acids SEQ ID NO 1 or 3 for increasing the starch or amylose content in plants, e.g. potato plants which, as wild type, are capable of producing starch or amylose, see examples 2-12.

- 10 The invention is not limited to the over-expression of the nucleic acid sequences SEQ ID NO 1 or SEQ ID NO 3 in plants especially potato plants.

- 15 The over-expression of both nucleic acid sequences SEQ ID NO 1 and 3 can be used for enhancing amylose biosynthesis. Constructs containing the nucleic acids SEQ ID NO 1 and SEQ ID NO 3 can also be used for increasing the starch content or the amylopectin content in plants. These constructs can be made on the same T-DNA driven by one promoter each. These constructs can also be made on the same T-DNA in tandem driven by the same promoter. These constructs can also be transformed using more than one construct, either at the same time (co-transformation) or in different transformation events.

- 20 The above-described proteins and nucleic acids may be used for producing starch or amylose in transgenic plants.

- 25 The transfer of foreign genes into the genome of an organism, in particular of a plant, is referred to as transformation.

- For this purpose, methods known per se for transforming plants and regenerating plants from plant tissues or plant cells can be used, in particular in plants, for transient or stable transformation, e.g. as described in example 2.

- 30 Suitable methods for the transformation of plants are the protoplast transformation by polyethylene glycol-induced DNA uptake, the biolistic method using the gene gun - also known as particle bombardment method, electroporation, the incubation of dry embryos in a DNA-containing solution, microinjection and the above-described Agrobacterium-mediated gene transfer. Said methods are described, for example, in B. Jenes et al.,
35 Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, edited by S.D. Kung and R. Wu, Academic Press (1993), 128-143 and in Potrykus, Annu. Rev. Plant Physiol. Plant Molec. Biol. 42 (1991), 205-225).

The construct to be expressed is preferably cloned into a vector which is suitable for transforming *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al., Nucl. Acids Res. 12 (1984), 8711) or preferably pSUN2 (WO 02/00900).

- 5 Accordingly, the invention furthermore relates to vectors containing the above-described nucleic acids, nucleic acid constructs or expression cassettes.

- 10 *Agrobacteria* which have been transformed with an expression cassette can be used in a known manner for the transformation of plants, for example by bathing injured leaves or leaf sections in an *agrobacteria* solution and then culturing them in suitable media.

Apart from in plants, the expression cassette may also be used for transforming bacteria, in particular cyanobacteria, mosses, yeasts, filamentous fungi and algae.

- 15 Genetically modified plants, also referred to as transgenic plants herein below, are preferably prepared by cloning the fused expression cassette which expresses an amylose biosynthesis enhancing protein into a vector, for example pBin19, which is suitable for transforming *Agrobacterium tumefaciens*.
- 20 *Agrobacteria* which have been transformed with such a vector may then be used in a known manner for the transformation of plants, in particular of crop plants, for example by bathing injured leaves or leaf sections in an *agrobacteria* solution and then culturing them in suitable media.
- 25 The transformation of plants by *agrobacteria* is described, inter alia, in F.F. White, Vectors for Gene Transfer in Higher Plants; in Transgenic Plants, Vol. 1, Engineering and Utilization, edited by S.D. Kung and R. Wu, Academic Press, 1993, pp. 15-38. Transgenic plants which contain a gene for expression of a nucleic acid encoding an amylose biosynthesis enhancing protein, which has been integrated into the expression
- 30 cassette, can be regenerated in a known manner from the transformed cells of the injured leaves or leaf sections.

- 35 A host plant is transformed with a nucleic acid SEQ ID NO 1 or 3 encoding an amylose biosynthesis enhancing protein by incorporating an expression cassette as insertion into a recombinant vector whose vector DNA comprises additional functional regulatory signals, for example sequences for replication or integration. Suitable vectors are described inter alia, in Methods in Plant Molecular Biology and Biotechnology (CRC Press), chapter 6/7, pp. 71-119 (1993).

By way of example, the plant expression cassette may be incorporated into a derivative of the transformation vector pBin-19 with 35s promoter (Bevan, M., Nucleic Acids Research 12: 8711-8721 (1984).

- 5 Using the above-cited recombination and cloning techniques, it is possible to clone the expression cassettes into suitable vectors for maintenance and propagation of genetic material for example in *E. coli*. Suitable cloning vectors are, inter alia, pBR322, pUC series, M13mp series and pACYC184. Particularly suitable are binary vectors which can replicate both in *E. coli* and in *agrobacteria*.

10

The invention therefore further relates to the use of the above-described nucleic acids or of the above-described nucleic acid constructs, in particular of the expression cassettes, for preparing genetically modified plants or for transforming plants, plant cells, plant tissues or parts of plants.

15

The use is preferably aimed at increasing the starch or amylose content of the plant, of the tubers or in other parts of the plant.

20

The use is most preferably aimed at increasing the starch or amylose content of wild-type or transgenic potato plants and especially the tubers of wild-type or transgenic potato plants.

25

Accordingly, the invention further relates to a method for preparing genetically modified plants by introducing an above-described nucleic acid or an above-described nucleic acid construct into the genome of the starting organism.

30

The invention further relates to the genetically modified organisms, the genetic modification increasing the activity of an amylose biosynthesis enhancing protein compared to a wild type or transgenic plant and the amylose biosynthesis enhancing protein comprising the amino acid sequence SEQ ID NO 2 or 4 or a sequence which is derived from this sequence by substitution, insertion or deletion of amino acids and which is at least 50% identical at the amino acid level to the sequence SEQ ID NO 2 or 4.

35

As illustrated above, the amylose biosynthesis enhancing protein activity is increased compared to the wild type or transgenic plant preferably by increasing the gene expression of a nucleic acid encoding an amylose biosynthesis enhancing protein.

40

In a further preferred embodiment, gene expression of a nucleic acid encoding an amylose biosynthesis enhancing protein is increased, as illustrated above, by introducing nucleic acids encoding an amylose biosynthesis enhancing protein into the organism

and thus by over-expressing nucleic acids encoding an amylose biosynthesis enhancing protein.

5 Such transgenic plants, their propagation material and their plant cells, plant tissues, plant parts or tubers are a further subject of the present invention.

10 Genetically modified plants of the invention, which have an increased starch or amylose content and which can be consumed by humans and animals, can also be used as food- or feedstuffs or as feed and food supplements, for example directly or after processing known per se. The genetically modified plants may furthermore be used for producing starch or amylose-containing extracts of said plant and/or for producing feed and food supplements.

The invention further relates to:

- 15
- I. A polynucleotide that encodes a polypeptide of SEQ ID NO 1 or 3.
 - II. A polynucleotide comprising at least 30 contiguous bases of SEQ ID NO 1 or 3.
 - 20 III. A polynucleotide having at least 60 % sequence identity to SEQ ID NO 1 or 3, wherein the identity is based on the entire coding sequence.
 - IV. A polynucleotide having at least 60 % sequence identity to SEQ ID NO 1 or 3, wherein the % sequence identity is based on the entire sequence.
 - 25 V. A polynucleotide which selectively hybridizes, under stringent conditions and a wash in 2 X SSC at 50 °C, to a hybridization probe derivable from the polynucleotide sequence as set forth in SEQ ID NO 1 or 3, or from the genomic sequence.
 - 30 VI. A polynucleotide complementary to a polynucleotide of V.
 - VII. The polynucleotide of I, wherein the starch or amylose biosynthesis enhancing polynucleotide is from *Solanum tuberosum*.
 - 35 VIII. The polynucleotide of I encoding a polypeptide, which after over-expression in a plant cell increases the starch or amylose content.
 - IX. The polynucleotide of I in antisense orientation, which after expression in a
40 plant cell decreases the starch or amylose content.

- X. A vector comprising at least one polynucleotide of I.
- 5 XI. An expression cassette comprising at least one polynucleotide of I operably linked to a promoter, wherein the polynucleotide is in sense or antisense orientation.
- XII. A host cell which is introduced with at least one expression cassette of X.
- 10 XIII. The host cell of XI that is a plant cell.
- XIV. A transgenic plant comprising at least one expression cassette of XI.
- XV. The transgenic plant of XIII, wherein the plant is *Solanum tuberosum*.
- 15 XVI. A tuber from the transgenic plant of XIV.
- XVII. An isolated protein comprising a member selected from the group consisting of:
- 20 a) a polypeptide comprising at least 10 contiguous amino acids of SEQ ID NO 2 or 4,
- b) a polypeptide which is a plant amylose biosynthesis enhancing protein,
- c) a polypeptide comprising at least 55 % sequence identity to SEQ ID NO 2 or 4, wherein the sequence identity is based on the entire sequence and has at least one epitope in common with an amylose biosynthesis enhancing protein.
- 25 d) a polypeptide encoded by a polynucleotide selected from SEQ ID NO 1 or 3,
- e) a polypeptide of SEQ ID NO 2 or 4.
- 30 XVIII. The protein of XVII, wherein the polypeptide is catalytically active.
- XIX. A ribonucleic acid sequence encoding the protein of XVIII.
- XX. A method for modulating the level of amylose biosynthesis enhancing protein in a plant, comprising:
- 35 a) stably transforming a plant cell with a polynucleotide coding for an amylose biosynthesis enhancing protein operably linked to a promoter, wherein the polynucleotide is in sense or antisense orientation
- b) growing the plant cell under plant growing conditions to produce a regenerated plant capable of expressing the polynucleotide for a time sufficient
- 40

to modulate the level of amylose biosynthesis enhancing protein in the plant.

5 XXI. The method of XX, wherein the polynucleotide coding for an amylose biosynthesis enhancing protein is selected from SEQ-ID NO 1 or 3.

XXII. The method of XX, wherein the plant is *Solanum tuberosum*.

10 XXIII. The method of XX, wherein activity of the amylose biosynthesis enhancing protein is increased.

XXIV. A method for modulating the level of starch or amylose in a plant, comprising:
15 a) stably transforming a plant cell with a polynucleotide coding for an amylose biosynthesis enhancing protein operably linked to a promoter, wherein the polynucleotide is in sense or anti-sense orientation,
b) growing the plant cell under plant growing conditions to produce a regenerated plant capable of expressing the polynucleotide for a time sufficient to modulate level of starch or amylose in the plant.

20 XXV. A method for modulating the level of amylose in a plant, comprising:
a) stably transforming a plant cell with a polynucleotide encoding an amylose biosynthesis enhancing protein operably linked to a promoter, wherein the polynucleotide is in sense or anti-sense orientation.
25 b) growing the plant cell under plant growing conditions to produce a regenerated plant capable of expressing the polynucleotide for a time sufficient to modulate level of amylose in the plant.

30 XXVI. The method of XXIV wherein the polynucleotide coding for an amylose biosynthesis enhancing protein is selected from SEQ ID NO 1 or 3.

Some of the terms used further on in the specification are defined at this point.

35 "Enzymatic activity/activity assay": the term enzymatic activity describes the ability of an enzyme to convert a substrate into a product. In this context, both the natural substrate of the enzyme and a synthetic modified analog of the natural substrate can be used. The enzymatic activity can be determined in what is known as an activity assay via the increase in the product, the decrease in the starting material, the decrease or increase in a specific cofactor, or a combination of at least two of the aforementioned parameters as a function of a defined period of time.
40

"Functional equivalents" in the present context describe nucleic acid sequences which hybridize under standard conditions with the nucleic acid sequence encoding the amylose biosynthesis enhancing protein or portions of the nucleic acid sequence encoding the amylose biosynthesis enhancing protein, and which are capable of bringing about the expression of an enzymatically active plant amylose biosynthesis enhancing protein in a cell or an organism.

It is advantageous to use short oligonucleotides of a length between 10 to 50bp, preferably 15-40bp, for example of the conserved or other regions, which can be determined via comparisons with other related genes in a manner known to the skilled worker for the hybridization. Alternatively, it is also possible to use longer fragments of the nucleic acids according to the invention or the complete sequences for the hybridization. These standard conditions vary depending on the nucleic acid used, namely oligonucleotide, longer fragment or complete sequence, or depending on which type of nucleic acid, that is DNA or RNA, is being used for the hybridization. Thus, for example, the melting temperatures for DNA:DNA hybrids are approx. 10°C lower than those of DNA:RNA hybrids of equal length. Suitable hybridization conditions are described above.

A functional equivalent is furthermore also understood as meaning, in particular, natural or artificial mutations of the relevant nucleic acid sequences of the plant amylose biosynthesis enhancing protein and their homologs from other organisms which make possible the expression of the enzymatically active plant amylose biosynthesis enhancing protein in a cell or an organism.

Thus, the scope of the present invention also extends to, for example, those nucleotide sequences which are obtained by modification of the nucleic acid sequence of a amylose biosynthesis enhancing protein. The purpose of such a modification can be, for example, the insertion of further cleavage sites for restriction enzymes, the removal of excess DNA, or the addition of further sequences. Proteins which are encoded via said nucleic acid sequences should still maintain the desired functions, despite the deviating nucleic acid sequence.

The term functional equivalent may also refer to the protein encoded by the nucleic acid sequence in question. In this case, the term functional equivalent describes a protein whose amino acid sequence is up to a specific percentage identical with that of the amylose biosynthesis enhancing protein.

Functional equivalents thus encompass naturally occurring variants of the sequences described herein, and also artificial, for example chemically synthesized, nucleic acid

sequences adapted to the codon usage, or the amino acid sequences derived therefrom.

5 In general, it can be said that functional equivalents independently of the amino acid sequence in question (encoded by a corresponding nucleic acid sequence) have in each case the enzymatic activity of a amylose biosynthesis enhancing protein.

10 "Reporter genes" encode readily quantifiable proteins. Using these genes, an assessment of transformation efficacy or of the site or time of expression can be made via growth, fluorescence, chemoluminescence, bioluminescence or resistance assay or via photometric measurement (intrinsic color) or enzyme activity. Very especially preferred in this context are reporter proteins (Schenborn E, Groskreutz D. Mol. Biotechnol. 1999; 13(1):29-44) such as the "green fluorescence protein" (GFP) (Gerdes HH and Kaether C, FEBS Lett. 1996; 389(1):44-47; Chui WL et al., Curr. Biol. 1996, 6:325-330; 15 Leffel SM et al., Biotechniques. 23(5):912-8, 1997), chloramphenicol acetyl transferase, a luciferase (Giacomin, Plant Sci. 1996, 116:59-72; Scikantha, J. Bact. 1996, 178:121; Millar et al., Plant Mol. Biol. Rep. 1992 10:324-414), and luciferase genes, in general b-galactosidase or b-glucuronidase (Jefferson et al., EMBO J. 1987, 6, 3901-3907), the Ura3 gene, the Ilv2 gene, the 2-desoxyglucose-6-phosphate phosphatase gene, b-lactamase gene, the neomycin phosphotransferase gene, the hygromycin phosphotransferase gene, or the BASTA (= gluphosinate) resistance gene. 20

25 "Significant increase": referring to the enzymatic activity, is understood as meaning the increase in the enzymatic activity of the enzyme incubated with a candidate compound in comparison with the activity of an enzyme not incubated with the candidate compound, which lies outside an error in measurement.

30 "Substrate": Substrate is the compound which is recognized by the enzyme in its original function and which is converted into a product by means of a reaction catalyzed by the enzyme.

Preferably, the plant amylose biosynthesis enhancing protein is encoded by a nucleic acid sequence comprising

- 35 a) a nucleic acid sequence shown in SEQ ID NO 1 or 3; or
- b) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from the amino acid sequence shown in SEQ ID NO 2 or 4 by back translation; or

- c) a nucleic acid sequence which, owing to the degeneracy of the genetic code, can be deduced from a functional equivalent of the amino acid sequence shown in SEQ ID NO 2 or 4, which has an identity with SEQ ID NO 2 or 4 of at least 50%; by back translation.

5

The functional equivalent of SEQ ID NO 2 or 4 set forth in c) has an identity of at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57% preferably at least 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, and 70% more preferably 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85% most preferably at least 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity with the SEQ ID NO 2 or 4.

10

Example 1.

15 Complementation study in yeast

Yeast contains two self-glycosylating proteins, Glg1p and Glg2p, which yield primers for the initiation of glycogen synthesis. For glycogen synthesis to take place in yeast it is required that either gene is functional. Yeast strain CC9, contain knock-out mutations for both genes and is therefore a null mutant regarding this specific biosynthetic function and is therefore unable to produce glycogen (Cheng, C. et al., Molecular and Cellular Biology (1995), 6632-6640). CC9 was used as a basis for complementation experiments with the isolated potato genes in order to validate their function by restoring glycogen biosynthesis in the CC9 strain. The potato genes were cloned in a yeast plasmid, pRS414 (Stratagene), and expressed with various yeast controlling elements such as Gal1, Adh1 and Glg2p promoters. CC9 was transformed by the resulting plasmids using LiCl and electroporation (Multiporator, Eppendorf). Transformed yeast colonies growing on appropriate media plates were screened by immersing in iodine solution. Wild type yeast producing glycogen is stained red brown by iodine while the null mutant CC9 is not stained. CC9 expressing the potato genes, StGH1 and StGH2, will stain red brown, when the isolated genes complement a glycogenin function in yeast and thus carry the desired function.

20

25

30

Example 2

35

Transformation method

Fully expanded leaves from in vitro propagated potato plants are diagonally cut in 2-4 pieces and precultivated on MC-plates for 2-3 days at 23-24°C.

40

Agrobacterium tumefaciens strain LBA4404 containing pHS1, pHS2, pHS3 or pHS4 are grown in YEB medium with 100 µg rifampicin and 25 µg/ml kanamycin over night on constant shaking (200 rpm) at 28°C.

- 5 The *Agrobacterium* culture is prepared for infection by dilution 1:20 with MS10 medium. The leaf explants are infected for 8-10 min in the bacterial solution and afterwards drained on filter paper for 5-20 seconds. The leaf segments are placed on the MS300 plates for 2 days co-cultivation under modest light at 23-24°C. At the end of co-cultivation the leaf segments are moved to M400 plates containing 400mg/l Claforan to suppress bacterial growth. After 4-5 days the explants are moved to selection medium
- 10 MS400 supplemented with 400mg/l Claforan. For explants transformed with pHS1 and pHS2 50 µM kanamycin was included in the media and for explants transformed with pHS3 and pHS4 0.5 µM Imazamox was added to the media. Leaf segments are transferred to fresh MS 400 selection medium every fortnight. The
- 15 regenerated putative transgenic shoots are collected and cultivated on MS30 plates with 200mg/l Claforan aiming at shoot elongation.

- 20 When the shoots are 3-5 cm long, 1-2 cm are cut off and grown on microtuber medium in the dark at 25°C. After 2-5 weeks microtubers are produced. Putative transgenic plants are analysed for GUS expression in microtubers to determine the transformation efficiency.

MC plates	MS300
MS300 plates with 1.5-2 ml liquid MS100 medium and covered with one sterile filter paper	4.4g/l MS-medium 2mg/l naphthyl acetic acid 1mg/l 6-benzyl amino pyridine 3% (w/v) sucrose pH 5.2
MS10	MS400
4.4 g/l MS-medium (murashige and Skoog) 1% (w/v) sucrose pH 5.8	4.4 g/l MS-medium 2mg/l zeatine 0.01mg/l naphthyl acetic acid mg/l gibberellic acid 10% (w/v) sucrose 400 mg/l claforan 0.5 µM Imazamox or 50 µM kanamycin pH5.8

PH 5,8	Mg/l gibberellic acid 10% (w/v) sucrose 400 mg/l claforan 0,5 µM Imazamox or 50 µM kanamycin pH 5.8
µM MS30 4.4 g/l MS-medium 3% (w/v) sucrose pH 5.8	Microtuber medium 4.4 g/l MS- medium 2.5 mg/l kinetin 0.5 mg/l abscisic acid 8% sucrose 200 mg/ claforan
MS100 4.4 mg/l MS-medium 30g/l sucrose 0.5 mg/l thiamin-HCl 0.5 mg/l pyridoxin-HCl 1 mg/l nicotinacid 0.5 mg/l kinetin 29.8 mg/l ferrous sulfate hepta hydrate 1 mg/l 2,4-Dichlorophenoxyacetic acid 2 g/l caseinhydrolysate pH 5.2	

Example 3

Transgenic plant AM 99-2003

5

High amylose potato lines can be produced for example by using antisense, RNAi or antibody technology that target the two starch branching enzymes starch branching enzyme 1 (SBE1) and starch branching enzyme 2 (SBE2).

- 10 The high amylose potato line AM99-2003 is produced by inhibition of the starch branching enzyme activities in the mother variety Dinamo. Transformation is made with a construct of SBE1 and SBE2 in antisense orientation driven by the gbss promoter.

- 15 pBluescript containing a 1620bp fragment of the 3' end of Sbe1 between EcoRV and SpeI is cut open with SpeI (blunt) and XbaI and ligated with a 1243bp SstI (blunt) and

XbaI fragment of the 3' end of Sbe2. The Sbe2 and Sbe1 complex is cut out with EcoRV and XbaI and ligated to the SmaI and XbaI opened up binary vector pHo3.1, see figure 8. The final vector is named pHAb12A, see figure 9 and nucleic acid sequence SEQ ID NO 15. pHo3.1 is based on pGPTVKan (Becker, D. et al., Plant Molecular Biology 20 (1992), 1195-1197) with the addition of the 987bp gbss promoter cloned at the HindIII site of pGPTVKan and the uidA gene is deleted by SmaI and SstI.

The mother variety Dinamo is transformed with the construct pHAb12A as described in example 2.

Example 4

Down-regulation of StGH1 and StGH2 genes in potato by antisense

The StGH1 and StGH2 genes were down-regulated in potato by transformation with the genes in antisense direction in relation to a plant regulatory element. The respective antisense genes were cloned in a binary vector driven by a tuber specific gbss promoter. NptII, yielding resistance to the antibiotic kanamycin, was used as selection marker. Two varieties were transformed, Prevalent and Producent. The shoots were selected on 50 µM kanamycin, which is a standard kanamycin concentration used for potato transformation (Ooms, G et al., Theoretical and Applied Genetics 73:744-750 (1987) and Tavazza, R. et al., Plant Science 59 (1988), 175-181).

Example 5

Over-expression of StGH1 and StGH2 genes in potato

The StGH1 and StGH2 genes were over-expressed in potato driven by the tuber specific promoter gbss. A mutated AHAS gene was used as selection marker yielding tolerance to the Imazamox herbicides. Two potato varieties were transformed, Desiree and AM99-2003 a transgenic high amylose line with a 40% decrease in starch content compared to its mother variety.

Example 6

Selection of transgenic lines

Non-transgenic escapes were identified and discarded by a PCR screening method. DNA was extracted according to DNeasy 96 Plant protocol (Qiagen). In a 96 well microtiter plate, 10-15 mg leaf tissue was added to each well together with a 5mm steel

ball, each well then representing one individual shoot. The plates were frozen in N₂(l) before homogenisation. The homogenisation was done at 30Hz in a Mixermill300 for 1 min. The DNA was at the end of the extraction protocol eluted in 75µl H₂O.

- 5 Specific primers for nptII and AHAS were used for the amplification of a 246bp fragment respective a fragment of 509 bp for selection of successfully transformed lines.

Npt2_for 5'-AGCAAGGTGAGATGACAGGAGATC-3'

Npt2_rev 5'CAGACAATCGGCTGCTCTGATG-3'

10

AHAS1_frw: 5'-AACAAACAACATCTTCTTCGATC-3'

AHAS1_rev: 5'-TAACGAGATTTGTAGCTCCG-3'.

The PCR reactions were with the extracted DNA setup and run as follow:

15

Reaction:

10x PCR Mix 2,0 µl

Primer frw (25µM) 0,4 µl

20 Primer rev (25µM) 0,4 µl

dNTPs (10mM) 0,4 µl

RedTAQ (Sigma) 1,0 µl

Templat (~20ng/µl) 4,0 µl

H₂O 11,8 µl

25

PCR program:

94°C 30 s

59°C 30 s x29 cycles

30 72°C 30s

72°C 7 min

8°C Hold

Example 7

35

Gene expression analyses

The gene expression levels of the StGH1 and StGH2 genes were analysed in the transgenic potato lines with real-time PCR (ABI prism 7900HT, Applied Biosystems).

40 With real-time PCR the change of gene expression can be analysed regarding RNA

expression levels. For pHS1 and pHS2 transgenic lines, expression of both sense and antisense RNA of StGH1 and StGH2 was measured, while in pHS3 and pHS4 transgenic lines the change in StGH1 and StGH2 mRNA expression was analysed.

- 5 The target for pHS1 and pHS2 is to reduce transcript levels of StGH1 and StGH2 respectively while the target for pHS3 and pHS4 is to increase transcript levels of the respective genes.

- 10 RNA was isolated from microtubers of the transgenic potato lines and mother varieties using Invisorb Spin Plant-RNA mini kit (Invitek). A reverse transcription reaction was made with 250 ng total RNA in 25 µl total reaction volume using TaqMan reverse transcription reagents (Applied Biosystems). Separate and specific primers (see table 1) were designed and used for the reverse transcription reaction in order to be able to differentiate the endogenous expression from the antisense RNA expression of the

15

StGH1 sense RNA	5'-TGAAGACAGCACAAAACCTGG-3'
StGH1 antisense RNA	5'-GTGAAAGTTTGAACGCACAC-3'
StGH2 sense RNA	5'-AGTGCCATAACATGCTTTCC-3'
StGH2 antisense RNA	5'-CACATTTTCAGCTGTTGATGGA-3'

Table 1

- 20 5 µl of the reverse transcription reaction was used in triplicate analyses together with specific sequence detection primers, TaqMan MGB probe (see table 2) and UMM mastermix (Applied Biosystems) and determined with real-time PCR.

StGH1	
Forward Primer:	TCGAGTCGCCACGTTAGAACTC
Reverse primer:	GAAATGCGTATGCGACTATGATG
TaqMan probe:	AGTCTCTCGGAGTTCC
StGH2	
Forward primer:	GGTGCTGATCCTCCAGTTCTCT
Reverse primer:	GTCCCTGAAGCATAACCAAGGT
TaqMan probe:	TTCTGCACTACTTAGGCCT

Table 2

Down-regulation of the two genes resulted in a decrease in gene expression in transgenic lines compared to their mother varieties in the order of 50-95 %.

- 5 Over-expression of the two genes resulted in a 2-10 times increase in gene expression in transgenic lines compared to their mother varieties.

Ref no	Construct	Variety	ΔCt s	Times increase or decrease in gene expression compared to mother variety
P01-041-84	pHS1	Producent	-1,14	-1,3
P02-325-1	pHS1	Producent	-2,03	-4,1
P02-325-9	pHS1	Producent	-1,47	-2,2
P02-325-11	pHS1	Producent	-1,25	-1,6
P02-325-15	pHS1	Producent	-2,52	-6,3
P02-325-25	pHS1	Producent	-2,0	-4
P02-325-27	pHS1	Producent	-1,64	-2,7
P02-325-33	pHS1	Producent	-1,59	-2,5
P02-325-34	pHS1	Producent	-1,52	-2,3
P02-325-63	pHS1	Producent	-1,53	-2,3
P02-300-37	pHS2	Prevalent	-1,27	-1,6
P02-300-66	pHS2	Prevalent	-1,04	-1,1
P02-300-71	pHS2	Prevalent	-1,13	-1,3
P02-300-73	pHS2	Prevalent	-1,1	-1,2
P02-300-80	pHS2	Prevalent	-2,12	-4,5
P02-300-127	pHS2	Prevalent	-1,67	-2,8
P02-300-140	pHS2	Prevalent	-3,96	-15,7
P02-303-31	pHS2	Prevalent	-1,16	-1,4
P02-303-64	pHS2	Prevalent	-1,15	-1,3
P02-305-54	pHS2	Prevalent	-1,33	-1,8
P02-320-24	pHS2	Prevalent	-1,03	-1,1
P02-307-4	pHS3	Desirée	1,82	3,3
P02-307-5	pHS3	Desirée	2,68	7,2
P02-307-12	pHS3	Desirée	2,67	7,1
P02-307-14	pHS3	Desirée	1,83	3,3
P02-307-15	pHS3	Desirée	1,79	3,2
P02-307-33	pHS3	Desirée	3,21	10,3
P02-307-43	pHS3	Desirée	2,7	7,3
P02-307-51	pHS3	Desirée	2,73	7,5
P02-307-80	pHS3	Desirée	2,78	7,7
P02-307-87	pHS3	Desirée	1,02	1,1

Ref no	Construct	Variety	ΔCt s	Times increase or decrease in gene expression compared to mother variety
P02-307-148	pHS3	Desirée	1,88	3,5
P02-309-63	pHS3	AM99-2003	1,64	2,7
P02-309-106	pHS3	AM99-2003	1,75	3,1
P02-311-59	pHS3	AM99-2003	1,17	1,4
P02-312-15	pHS4	AM99-2003	1,03	1,1
P02-313-21	pHS4	AM99-2003	1,54	2,4

Table 3: Gene expression analyses based on Real- Time PCR

Example 8

5

Dry matter analysis

Dry matter has been analysed on microtubers from transgenic lines transformed with pHS1, pHS2, pHS3 and pHS4 showing a down-regulation or over expression of the genes. Since starch normally contribute to more than 80% of the dry matter in potato tubers, an increase or decrease in starch content will affect also the dry matter content.

10

Two microtubers of each line were harvested when they had reached maturity. Dry matter was calculated for mature microtubers weighed before and after 72 hours drying at 60°C. For comparison microtubers from the varieties Dinamo, Desirée, Prevalent, Producent and P737 with starch contents between 13 and 28% (when grown in field) were used. The starch content of microtubers is not as high as starch content of field grown tubers. However dry matter content can readily be compared in microtubers and that value is well correlated to the determined starch content in field grown tubers. In table 4 the average dry matter for the different varieties, calculated on ten or more microtubers, is shown.

15

20

Variety	Starch content field grown tubers	Dry matter microtubers
Dinamo	13%	14,8
Desirée	16%	16,1
Producent	22%	19,2
Prevalent	22%	19,7
P737	28%	21,6

Table 4. Dry matter content of 5 varieties based on 10 or more microtubers

One of each pHS1 and pHS2 with confirmed decrease in gene-expression have been analyzed for dry matter so far. Those two have a decrease in dry matter of 7 and 11% compared to their mother varieties.

- 5 For the pHS3 lines 8 of 9 of the confirmed over-expressed lines show an increase of up to 36% in dry matter. See table 5.

Ref No	Construct	Variety	Dry matter in relation to mother variety (%)
41-84	pHS1	Producent	89
300-127	pHS2	Prevalent	93
300-140	PHS2	Prevalent	96
307-4	pHS3	Desirée	106
307-5	pHS3	Desirée	117
307-15	pHS3	Desirée	124
307-33	pHS3	Desirée	116
307-57	pHS3	Desirée	136
309-63	pHS3	AM99-2003	134
309-106	pHS3	AM99-2003	109
309-111	pHS3	AM99-2003	108

Table 5. Dry matter content on transgenic lines with confirmed down-regulation or over expression of the StGH1 and StGH2 genes

Example 9

Starch content analysis

For analysis of starch content a total starch assay procedure from Megazyme International Ireland Ltd., Bray, Co. Wicklow, Ireland (AOAC Method 996.1; AACC method 76.13; ICC standard method No. 168) was used according to the suppliers instructions.

- 20 Starch content was analysed on microtubers from all transgenic lines transformed with pHS1, pHS2, pHS3 and pHS4. The microtubers were harvested when they had reached maturity. Mature microtubers were ground and maltosaccharides and free glucose residues were washed away with ethanol. The microtuber starch was treated with DMSO to ensure the complete solubilisation of samples with high levels of resistant starch, as the high amylose clones.

Samples were analysed with a standard spectrophotometric assay procedure. The transgenic lines were compared to potato varieties with known starch content ranging from 8% to 30%. The results give an indication on the change in starch content related to the genetic modification of the different transgenic lines.

5

Example 10

Field-testing of transgenic potato lines

10 Produced transgenic lines are tested in field trials for determination of agronomic performance in relation to mother variety and other varieties used for starch production. Starch content, which is a main agronomic factor of importance for crops used for starch processing, can be measured by several different methods.

15 Under water weighing is performed on a scale in a tub of water and where an increase in starch content is associated with an increase in density of the sample. An increase in starch is associated with an increased dry matter content, which can be measured by comparing the tissue fresh weight to tissue dry weight after extensive water elimination in an oven.

20

Starch content can also be measured by enzymatic methods as described under starch content analyses.

Example 11

25

Increased amylose and starch yield in potatoes

Potato varieties used for starch production as well as genotypes with a high amylose content are transformed with gene constructs as described above for the over-expression of a starch biosynthesis enhancing protein. The over-expression of StGH1 or StGH2 in potato plants will result in an increased amylose content of the transgenic plant compared to the starting plant.

30

Example 12

35

Increased solids and improved processing quality of potatoes

In another aspect the invention may be used to increase the solids content of varieties that are used for processed potato products or table potato varieties. The potato genotypes are transformed with gene constructs as described above for the over-expression

40

of a starch biosynthesis enhancing protein. This starch biosynthesis enhancing protein may be derived from genes described above or other plant genes containing the same functional domains.

What is claimed:

1. A method of increasing the production of starch in plants comprising culturing a plant with enhanced expression or activity of at least one starch biosynthesis enhancing protein;
2. The method of claim 1, wherein said starch has a high amylose content.
3. The method as claimed in either claim 1 or 2, wherein production of amylose is increased.
4. The method as claimed in any one of claims 1 to 3, wherein said method comprises over-expression of an amylose biosynthesis enhancing protein.
5. The method as claimed in claim 4, wherein said protein comprises the SEQ ID NO: 2 or 4 or a protein derived from this sequence by substitution, insertion or deletion of amino acids and which has at least 50% identity at the amino acid level with SEQ ID NO: 2 or 4.
6. The method as claimed in any of claims 1 to 5, wherein the amylose biosynthesis enhancing protein is encoded by a nucleic acid sequence selected from the group consisting of:
 - a) a nucleic acid sequence comprising a nucleotide sequence which is at least 60% identical to the nucleic acid sequence of SEQ ID NO: 1 or 3;
 - b) a nucleic acid sequence comprising a fragment of at least 30 nucleotides of a nucleic acid sequence comprising the nucleotide sequence of SEQ ID NO: 1 or 3;
 - c) a nucleic acid sequence which encodes a polypeptide comprising an amino acid sequence at least about 60% identical to the amino acid sequence of SEQ ID NO: 2 or 4 and
 - d) a nucleic acid sequence which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 or 4 or wherein the fragment comprises at least 10 contiguous amino acid residues of the amino acid sequence of SEQ ID NO: 2 or 4,

7. The method as claimed in any one of claims 1 to 6, wherein the amylose biosynthesis enhancing protein is encoded by a nucleic acid sequence comprising the nucleotide sequence set forth as SEQ ID NO:1 or SEQ ID NO:3.
- 5 8. The method as claimed in any one of claims 1 to 7, wherein deficiency or decreased activity is achieved by a method selected from the group consisting of:
- 10 a) knock-out of the gene encoding said protein;
- b) mutagenesis of the gene encoding said protein, wherein said mutation can be induced in the coding, non-coding, or regulatory regions of said gene;
- 15 c) expression of an anti-sense RNA, wherein said anti-sense RNA is complementary to at least part of the RNA encoding said protein;
9. A method of producing amylose type starch by culturing a plant which over-expresses SEQ ID NO:1 or 3 or has increased amylose biosynthesis enhancing activity under conditions such that the plant produces an increased amount of amylose type starch.
- 20 10. The method of any of the preceding claims, wherein said plant belongs to the genus *Solanum*.
- 25 11. The method of claim 10, wherein said plant is *Solanum tuberosum*.
12. A nucleic acid sequence SEQ ID NO:1 encoding an amylose biosynthesis enhancing protein.
- 30 13. A nucleic acid sequence SEQ ID NO:3 encoding an amylose biosynthesis enhancing protein.
14. An amino acid sequence SEQ ID NO:2 having amylose biosynthesis enhancing activity.
- 35 15. An amino acid sequence SEQ ID NO:4 having amylose biosynthesis enhancing activity.

16. A transgenic expression cassette comprising in combination with a regulatory sequence a nucleic acid sequence selected from the group consisting of:

- 5
- a) a nucleic acid sequence comprising a nucleotide sequence which is at least 60% identical to the nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3,
- b) a nucleic acid sequence comprising a fragment of at least 30 nucleotides of a nucleic acid sequence comprising the nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3,
- 10
- c) a nucleic acid sequence which encodes a polypeptide comprising an amino acid sequence at least about 60% identical to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4, or
- d) a nucleic acid sequence which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4
- 15
- wherein the fragment comprises at least 10 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4

wherein said regulatory sequence is capable of mediating expression of said nucleic acid sequence in a plant.

20

17. A transgenic expression cassette of claim 16, wherein said regulatory sequence is a promoter sequence heterologous with regard to said nucleic acid sequence.

25

18. A transgenic expression cassette of claim 16, wherein said regulatory sequence is a tuber specific promoter sequence.

30

19. A transgenic expression cassette of either claim 16, 17 or 18, wherein said nucleic acid sequence is arranged in antisense or sense orientation with regard to said promoter sequence.

20. A transgenic expression cassette of any of the claims 16 to 19, wherein said nucleic acid sequence encodes a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:4.

35

21. A transgenic expression cassette of any of the claims 16 to 20, wherein said nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO:1 or SEQ ID NO:3.

22. A transgenic expression cassette of any of the claims 16 to 21, wherein said nucleic acid sequence encodes a naturally occurring variant of a polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:4.
23. A transgenic host cell transformed with an expression cassette of any of the claims 16 to 22.
24. A transgenic host cell of claim 23, wherein said host cell belongs to the genus *Solanum*.
25. A transgenic plant comprising an expression cassette of any of claims 16 to 22.
26. A transgenic potato plant comprising an expression cassette of any of claims 16 to 22.
27. A transgenic potato plant, plant part, seed or tuber comprising an expression cassette of any of claims 16 to 22.

SEQUENCE LISTING

<110> BASF Plant Science GmbH

5 <120> Enhanced Amylose Production in Plants

<130> AE884-02

<140> PF0000054331

10 <141> 2003-03-07

<160> 15

15 <170> PatentIn Ver. 2.1

<210> 1

<211> 2084

<212> DNA

20 <213> Solanum tuberosum

<220>

<221> CDS

<222> (302)..(1696)

25 <400> 1

ttttcataaaa ctttttcaac tttattccat actcttttat ttatcagctc ctagatcttc 60

ttttttgttt gttgatattc tcttgaaaat tgttcagtga agagttgatc aaagctaaga 120

30 cacaggggct gcggccattt tttcacggga atctttcttct ttattttccg gtgaaagtct 180

gaacgcacac cgttatttct agacagtaga caatgtcaag tgaanaacat cacaagtctt 240

35 tgaagatttg taattaatta gtcgagattt ttaatttgga ggaaagagaa aaacagagaa 300

g atg ata ggg cgg gtg ggc ttg ttg ttg gta ttg ttg ata gca acg acg 349
Met Ile Gly Arg Val Gly Leu Leu Leu Val Leu Leu Ile Ala Thr Thr
1 5 10 1540 gtg act att ggg gct gaa acg acg acg tta aaa ggg gta aac aga aat 397
Val Thr Ile Gly Ala Glu Thr Thr Thr Leu Lys Gly Val Asn Arg Asn
20 25 3045 gcg tat gcg act atg atg tat atg gga act ccg aga gac tac gag ttc 445
Ala Tyr Ala Thr Met Met Tyr Met Gly Thr Pro Arg Asp Tyr Glu Phe
35 40 4550 tac gtg gcg act cga gta atg ctc cga tca ctt acc cgg cta gga gtt 493
Tyr Val Ala Thr Arg Val Met Leu Arg Ser Leu Thr Arg Leu Gly Val
50 55 6055 gaa gcc gat ctc gtc gtt att gct tca ctt gac gtt cct ctt cgc tgg 541
Glu Ala Asp Leu Val Val Ile Ala Ser Leu Asp Val Pro Leu Arg Trp
65 70 75 80gtt caa act cta gaa cag gaa gat ggt gct aag gtg gtg aga gtt aaa 589
Val Gln Thr Leu Glu Gln Glu Asp Gly Ala Lys Val Val Arg Val Lys
85 90 9560 aat ctg aac aat ccg tat tgt atc aac cct aat tgg aga ttc aag ctc 637
Asn Leu Asn Asn Pro Tyr Cys-Ile Asn Pro Asn Trp Arg Phe Lys-Leu
100 105 110

2

	aca ctg aac aaa ctt tat gcg tgg agc ctc gta aat tat gac agg gtt Thr Leu Asn Lys Leu Tyr Ala Trp Ser Leu Val Asn Tyr Asp Arg Val	685
	115 120 125	
5	gtc atg ctt gat gct gac aac ctt ttc ctc cag aaa act gat gaa ctg Val Met Leu Asp Ala Asp Asn Leu Phe Leu Gln Lys Thr Asp Glu Leu	733
	130 135 140	
10	ttc caa tgt ggc cag ttt tgt gct gtc ttc att aat ccc tgc atc ttc Phe Gln Cys Gly Gln Phe Cys Ala Val Phe Ile Asn Pro Cys Ile Phe	781
	145 150 155 160	
15	cac act ggt ctc ttt gta ttg cag cca tca aaa aag gtg ttc aat gac His Thr Gly Leu Phe Val Leu Gln Pro Ser Lys Lys Val Phe Asn Asp	829
	165 170 175	
20	atg atc cat gag ata gag att ggg agg gaa aat caa gac ggt gca gac Met Ile His Glu Ile Glu Ile Gly Arg Glu Asn Gln Asp Gly Ala Asp	877
	180 185 190	
	caa ggt ttc att gga ggc cac ttc cca gat tta ctt gat cgg cca atg Gln Gly Phe Ile Gly Gly His Phe Pro Asp Leu Leu Asp Arg Pro Met	925
	195 200 205	
25	ttc cac cct cct ctt aat ggt acc cag ctc cag gga agt tac agg ctt Phe His Pro Pro Leu Asn Gly Thr Gln Leu Gln Gly Ser Tyr Arg Leu	973
	210 215 220	
30	cct cta gga tac caa atg gac gcc tct tat tat tat ctc aaa ctc cat Pro Leu Gly Tyr Gln Met Asp Ala Ser Tyr Tyr Tyr Leu Lys Leu His	1021
	225 230 235 240	
35	tgg tgg gta cct tgt gga cct aat agt gtc att aca ttt cct ggt gct Trp Ser Val Pro Cys Gly Pro Asn Ser Val Ile Thr Phe Pro Gly Ala	1069
	245 250 255	
40	cca tgg tta aaa cca tgg tat tgg tgg tca tgg cct gtc tta ccc ttg Pro Trp Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro Leu	1117
	260 265 270	
	ggc atc cag tgg cat gaa cag cga cgt cta act gtt ggg tat ggt gct Gly Ile Gln Trp His Glu Gln Arg Arg Leu Thr Val Gly Tyr Gly Ala	1165
	275 280 285	
45	gag atg ata gca gtg ttg atc caa tct ata ttt tac cta gga ata att Glu Met Ile Ala Val Leu Ile Gln Ser Ile Phe Tyr Leu Gly Ile Ile	1213
	290 295 300	
50	gca gtg aca cgc cta gca cgc cca aat tta tca aag ttg tgc tat cgc Ala Val Thr Arg Leu Ala Arg Pro Asn Leu Ser Lys Leu Cys Tyr Arg	1261
	305 310 315 320	
55	cat gat gat agc aag agt gcc ttc tta cta cga act ggc ctt aaa ttg His Asp Asp Ser Lys Ser Ala Phe Leu Leu Arg Thr Gly Leu Lys Leu	1309
	325 330 335	
60	att gct ata tgg tcc att ott gct gcc tac aca gtt cct tat ttc gtg Ile Ala Ile Trp Ser Ile Leu Ala Ala Tyr Thr Val Pro Tyr Phe Val	1357
	340 345 350	
	att cct tgt aca gtt cat cca cta gtt ggc tgg agt ctc tac tta ctc Ile Pro Cys Thr Val His Pro Leu Val Gly Trp Ser Leu Tyr Leu Leu	1405

3

355 360 365

5 ggc tct ttt tca cta tcc tgt ata aca gtg aat gca ttt ctt ttg ccg 1453
 Gly Ser Phe Ser Leu Ser Cys Ile Thr Val Asn Ala Phe Leu Leu Pro
 370 375 380

10 atg cta cct gtt tta gto cca tgg att ggg atc ctt ggg gcc ctt ttg 1501
 Met Leu Pro Val Leu Val Pro Trp Ile Gly Ile Leu Gly Ala Leu Leu
 385 390 395 400

15 gtg atg gct tac cct tgg tac aac gac ggt gtt gta aga gca atg gct 1549
 Val Met Ala Tyr Pro Trp Tyr Asn Asp Gly Val Val Arg Ala Met Ala
 405 410 415

20 gta ttt aca tac gcc ttc tgt gct tct cca gca tta tgg atg gca ttg 1597
 Val Phe Thr Tyr Ala Phe Cys Ala Ser Pro Ala Leu Trp Met Ala Leu
 420 425 430

25 gtt aaa atc aag tgt tct ctt cat gtt tca ctt gag agg gaa gga ttc 1645
 Val Lys Ile Lys Cys Ser Leu His Val Ser Leu Glu Arg Glu Gly Phe
 435 440 445

30 ttg ccc aag ata agt gaa tct aca gca cct gct ggt tct aac aaa ctg 1693
 Leu Pro Lys Ile Ser Glu Ser Thr Ala Pro Ala Gly Ser Asn Lys Leu
 450 455 460

35 tat tgaaagtga aaagttaaag gaatcaacag gagaactaat gcttcagaaa 1746
 Tyr
 465

40 catctccaaa cgttttgctt aggagacttg gactctgctt gtgctatcct agctagttgc 1806
 ttacgtctgt gctcttaatt agaattggaat totgtgagtg ggtttagaat tgggaggatg 1866
 35 ttttgtgttg tacatggact atctotggtc tcttgaatgc tactccagga aaaagattgt 1926
 ttctcactta attttttctg ttactaaatt gtatgtggaa taggttcttt aaaattttatt 1986
 catggattta tgttatgtat gctaacagtg taaatattaa gtcttgggtga aataagtaat 2046
 40 tccttattca tacaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2084

45 <210> 2
 <211> 465
 <212> PRT
 <213> Solanum tuberosum

50 <400> 2
 Met Ile Gly Arg Val Gly Leu Leu Leu Val Leu Leu Ile Ala Thr Thr
 1 5 10 15
 Val Thr Ile Gly Ala Glu Thr Thr Thr Leu Lys Gly Val Asn Arg Asn
 20 25 30
 55 Ala Tyr Ala Thr Met Met Tyr Met Gly Thr Pro Arg Asp Tyr Glu Phe
 35 40 45
 60 Tyr Val Ala Thr Arg Val Met Leu Arg Ser Leu Thr Arg Leu Gly Val
 50 55 60
 Glu Ala Asp Leu Val Val Ile Ala Ser Leu Asp Val Pro Leu Arg Trp

	65		70		75		80									
	Val	Gln	Thr	Leu	Glu	Gln	Glu	Asp	Gly	Ala	Lys	Val	Val	Arg	Val	Lys
					85					90						95
5	Asn	Leu	Asn	Asn	Pro	Tyr	Cys	Ile	Asn	Pro	Asn	Trp	Arg	Phe	Lys	Leu
				100					105					110		
10	Thr	Leu	Asn	Lys	Leu	Tyr	Ala	Trp	Ser	Leu	Val	Asn	Tyr	Asp	Arg	Val
			115					120					125			
	Val	Met	Leu	Asp	Ala	Asp	Asn	Leu	Phe	Leu	Gln	Lys	Thr	Asp	Glu	Leu
		130					135					140				
15	Phe	Gln	Cys	Gly	Gln	Phe	Cys	Ala	Val	Phe	Ile	Asn	Pro	Cys	Ile	Phe
	145					150					155					160
	His	Thr	Gly	Leu	Phe	Val	Leu	Gln	Pro	Ser	Lys	Lys	Val	Phe	Asn	Asp
20					165					170					175	
	Met	Ile	His	Glu	Ile	Glu	Ile	Gly	Arg	Glu	Asn	Gln	Asp	Gly	Ala	Asp
			180					185					190			
25	Gln	Gly	Phe	Ile	Gly	Gly	His	Phe	Pro	Asp	Leu	Leu	Asp	Arg	Pro	Met
			195				200						205			
	Phe	His	Pro	Pro	Leu	Asn	Gly	Thr	Gln	Leu	Gln	Gly	Ser	Tyr	Arg	Leu
	210					215						220				
30	Pro	Leu	Gly	Tyr	Gln	Met	Asp	Ala	Ser	Tyr	Tyr	Tyr	Leu	Lys	Leu	His
	225				230					235						240
	Trp	Ser	Val	Pro	Cys	Gly	Pro	Asn	Ser	Val	Ile	Thr	Phe	Pro	Gly	Ala
35				245						250					255	
	Pro	Trp	Leu	Lys	Pro	Trp	Tyr	Trp	Trp	Ser	Trp	Pro	Val	Leu	Pro	Leu
			260					265					270			
40	Gly	Ile	Gln	Trp	His	Glu	Gln	Arg	Arg	Leu	Thr	Val	Gly	Tyr	Gly	Ala
		275						280					285			
	Glu	Met	Ile	Ala	Val	Leu	Ile	Gln	Ser	Ile	Phe	Tyr	Leu	Gly	Ile	Ile
	290					295					300					
45	Ala	Val	Thr	Arg	Leu	Ala	Arg	Pro	Asn	Leu	Ser	Lys	Leu	Cys	Tyr	Arg
	305				310						315					320
	His	Asp	Asp	Ser	Lys	Ser	Ala	Phe	Leu	Leu	Arg	Thr	Gly	Leu	Lys	Leu
50				325						330				335		
	Ile	Ala	Ile	Trp	Ser	Ile	Leu	Ala	Ala	Tyr	Thr	Val	Pro	Tyr	Phe	Val
			340					345					350			
55	Ile	Pro	Cys													

5

Val Met Ala Tyr Pro Trp Tyr Asn Asp Gly Val Val Arg Ala Met Ala
405 410 415

5 Val Phe Thr Tyr Ala Phe Cys Ala Ser Pro Ala Leu Trp Met Ala Leu
420 425 430

Val Lys Ile Lys Cys Ser Leu His Val Ser Leu Glu Arg Glu Gly Phe
435 440 445

10 Leu Pro Lys Ile Ser Glu Ser Thr Ala Pro Ala Gly Ser Asn Lys Leu
450 455 460

Tyr
465

15

<210> 3

<211> 2230

20

<212> DNA

<213> Solanum tuberosum

<220>

<221> CDS

25

<222> (143) .. (2086)

<400> 3

tatccccaga gaatcagctg aatcaagaac tgatttttag attatgtttt cttgattctt 60

30

tgaaatggga acttgatttt cagtttttca actcagatgt tgtgttcctt tagctggaaa 120

acttgaaaaa ggaaagccca ga atg aga gga agt tta got ggt gga cca cct 172
Met Arg Gly Ser Leu Ala Gly Gly Pro Pro
1 5 10

35

agt cct att gaa cct aga cag agg ctt tct gta ttc act gag gaa aca 220
Ser Pro Ile Glu Pro Arg Gln Arg Leu Ser Val Phe Thr Glu Glu Thr
15 20 25

40

agc aaa aga agg ttc ttg aga agt aaa gtt ttc aga gat ggg gag aga 268
Ser Lys Arg Arg Phe Leu Arg Ser Lys Val Phe Arg Asp Gly Glu Arg
30 35 40

45

gct ctt cat agt ccc acc aaa aac agg aat ttt acc tgc aag ttc cca 316
Ala Leu His Ser Pro Thr Lys Asn Arg Asn Phe Thr Cys Lys Phe Pro
45 50 55

50

act gtg aag ctt ata ttg ggt gtt att gct ctg gtt gca att tgg tca 364
Thr Val Lys Leu Ile Leu Gly Val Ile Ala Leu Val Ala Ile Trp Ser
60 65 70

55

ctc tgg cat tct cca gca att tat aac acg gaa tac ata tct agt tca 412
Leu Trp His Ser Pro Ala Ile Tyr Asn Thr Glu Tyr Ile Ser Ser Ser
75 80 85 90

ggc tot cgg gct gct ttg atg cac aga gag tta agt ggt cat tct tca 460
Gly Ser Arg Ala Ala Leu Met His Arg Glu Leu Ser Gly His Ser Ser
95 100 105

60

gct gat caa cgt tat aca tca ctt tta gat att gac tgg gac caa att 508
Ala Asp Gln Arg Tyr Thr Ser Leu Leu Asp Ile Asp Trp Asp Gln Ile
110 115 120

5	tcc caa gtt att gag aaa ctg gcc gat agg cat gag tat cag ggc gta	556
	Ser Gln Val Ile Glu Lys Leu Ala Asp Arg His Glu Tyr Gln Gly Val	
	125 130 135	
	ggg ata tta aac ttc aat gac agt gaa att gat cag ttg aag gag tta	604
	Gly Ile Leu Asn Phe Asn Asp Ser Glu Ile Asp Gln Leu Lys Glu Leu	
10	cta ccg gac gct gag cat gta atc ttg aac ctg gat cac gtc ccg aat	652
	Leu Pro Asp Ala Glu His Val Ile Leu Asn Leu Asp His Val Pro Asn	
	155 160 165 170	
	aat ata aca tgg gaa aca ata tat cct gaa tgg ata gat gaa gaa gaa	700
	Asn Ile Thr Trp Glu Thr Ile Tyr Pro Glu Trp Ile Asp Glu Glu Glu	
20	gaa ttt gag gtc ccc act tgt cct tct ctg ccc aaa att cag ttt ccg	748
	Glu Phe Glu Val Pro Thr Cys Pro Ser Leu Pro Lys Ile Gln Phe Pro	
	190 195 200	
	ggt aaa cca agg att gat ctc ata gtt gta aag ctt cca tgc aag aag	796
	Gly Lys Pro Arg Ile Asp Leu Ile Val Val Lys Leu Pro Cys Lys Lys	
25	tct aag gac tgg tat aga gat gta gct cgt ttt cac ttg cag ctg gca	844
	Ser Lys Asp Trp Tyr Arg Asp Val Ala Arg Phe His Leu Gln Leu Ala	
	220 225 230	
	gca gca agg ctg gct gcc agt aat aaa ggg tat cat cca ata cat gtg	892
	Ala Ala Arg Leu Ala Ala Ser Asn Lys Gly Tyr His Pro Ile His Val	
35	ctt ctg gtt act gag cat ttc cca acc ccc aat ctg ttc acc tgt aaa	940
	Leu Leu Val Thr Glu His Phe Pro Thr Pro Asn Leu Phe Thr Cys Lys	
	255 260 265	
	gag tta gtt gta cgt gaa ggc aat gca tgg cta tat gaa cct aat ctg	988
	Glu Leu Val Val Arg Glu Gly Asn Ala Trp Leu Tyr Glu Pro Asn Leu	
45	aac act tta aga gag aag ctc cac ctc cct gtt ggg tca tgt gaa ctt	1036
	Asn Thr Leu Arg Glu Lys Leu His Leu Pro Val Gly Ser Cys Glu Leu	
	285 290 295	
	gca gtt cct ctc aag gct aaa gca aat tgg cac tct gga aat gta aga	1084
	Ala Val Pro Leu Lys Ala Lys Ala Asn Trp His Ser Gly Asn Val Arg	
50	cga gaa gcc tat gca act att ctc cac tca gca aat ttt tat gta tgt	1132
	Arg Glu Ala Tyr Ala Thr Ile Leu His Ser Ala Asn Phe Tyr Val Cys	
	315 320 325 330	
	gga gcc ata gct gca gca cag agt att cgc ttg gca ggt tca acc cga	1180
	Gly Ala Ile Ala Ala Gln Ser Ile Arg Leu Ala Gly Ser Thr Arg	
60	gat ctt gtg ata ctt gtt gat gag act atc agt gac tac cac agg ggt	1228
	Asp Leu Val Ile Leu Val Asp Glu Thr Ile Ser Asp Tyr His Arg Gly	
	350 355 360	
	ggt tta gag gct gcc gga tgg aag atc cac acg ata aag aga ata agg	1276

7

	Gly	Leu	Glu	Ala	Ala	Gly	Trp	Lys	Ile	His	Thr	Ile	Lys	Arg	Ile	Arg	
	365							370					375				
5	aat	cct	aaa	gct	gaa	cag	gat	gcc	tac	aat	gag	tgg	aac	tat	agc	aaa	1324
	Asn	Pro	Lys	Ala	Glu	Gln	Asp	Ala	Tyr	Asn	Glu	Trp	Asn	Tyr	Ser	Lys	
	380						385				390						
10	ttt	cgt	ctc	tgg	cag	ctg	aca	gat	tat	gac	aaa	atc	atc	ttc	att	gat	1372
	Phe	Arg	Leu	Trp	Gln	Leu	Thr	Asp	Tyr	Asp	Lys	Ile	Ile	Phe	Ile	Asp	
	395				400					405					410		
15	gcg	gat	ttg	ttg	ata	ctg	aga	aat	att	gat	ttt	ctc	ttt	gag	atg	cct	1420
	Ala	Asp	Leu	Leu	Ile	Leu	Arg	Asn	Ile	Asp	Phe	Leu	Phe	Glu	Met	Pro	
					415					420					425		
	gaa	ata	act	gca	ata	gga	aat	aat	gca	acc	ctt	ttt	aat	tca	ggc	gtg	1468
	Glu	Ile	Thr	Ala	Ile	Gly	Asn	Asn	Ala	Thr	Leu	Phe	Asn	Ser	Gly	Val	
				430					435					440			
20	atg	gtc	gtt	gaa	cca	tca	aat	tgc	aca	ttt	cag	ctg	ttg	atg	gat	cat	1516
	Met	Val	Val	Glu	Pro	Ser	Asn	Cys	Thr	Phe	Gln	Leu	Leu	Met	Asp	His	
				445				450						455			
25	atc	aat	gag	att	gaa	tca	tac	aat	ggt	ggg	gat	cag	ggg	tat	ttg	aac	1564
	Ile	Asn	Glu	Ile	Glu	Ser	Tyr	Asn	Gly	Gly	Asp	Gln	Gly	Tyr	Leu	Asn	
		460					465				470						
30	gaa	ata	ttc	acc	tgg	tgg	cat	agg	atc	cca	aaa	cac	atg	aac	ttt	ttg	1612
	Glu	Ile	Phe	Thr	Trp	Trp	His	Arg	Ile	Pro	Lys	His	Met	Asn	Phe	Leu	
	475					480					485				490		
35	aaa	cat	tat	tgg	gaa	ggt	gat	gag	gag	gag	aag	aag	caa	atg	aaa	aca	1660
	Lys	His	Tyr	Trp	Glu	Gly	Asp	Glu	Glu	Glu	Lys	Lys	Gln	Met	Lys	Thr	
				495						500					505		
	cgt	ctt	ttt	ggt	gct	gat	cct	cca	gtt	ctc	tat	gtt	ctg	cac	tac	tta	1708
	Arg	Leu	Phe	Gly	Ala	Asp	Pro	Pro	Val	Leu	Tyr	Val	Leu	His	Tyr	Leu	
				510					515					520			
40	ggc	ctg	aaa	cct	tgg	tta	tgc	ttc	agg	gac	tac	gat	tgc	aac	tgg	aat	1756
	Gly	Leu	Lys	Pro	Trp	Leu	Cys	Phe	Arg	Asp	Tyr	Asp	Cys	Asn	Trp	Asn	
			525					530					535				
45	gtg	ggt	aag	ttg	cag	gag	ttt	gca	agt	gat	gtg	gca	cac	agg	acg	tgg	1804
	Val	Gly	Lys	Leu	Gln	Glu	Phe	Ala	Ser	Asp	Val	Ala	His	Arg	Thr	Trp	
		540					545					550					
50	tgg	aag	gtc	cat	gat	gcc	atg	cca	gat	aac	tta	cat	aaa	tat	tgt	ttg	1852
	Trp	Lys	Val	His	Asp	Ala	Met	Pro	Asp	Asn	Leu	His	Lys	Tyr	Cys	Leu	
	555					560					565				570		
55	ctt	agg	tct	aaa	cag	aag	gct	gca	cta	gag	tgg	gat	cga	aga	gaa	gct	1900
	Leu	Arg	Ser	Lys	Gln	Lys	Ala	Ala	Leu	Glu	Trp	Asp	Arg	Arg	Glu	Ala	
				575						580					585		
	gag	aaa	gct	aac	ttt	tca	gat	ggt	cat	tgg	aag	atc	aaa	ata	aag	gac	1948
	Glu	Lys	Ala	Asn	Phe	Ser	Asp	Gly	His	Trp	Lys	Ile	Lys	Ile	Lys	Asp	
				590					595					600			
60	cca	cgt	ttg	gag	act	tgt	tat	gaa	gaa	ttt	tgc	ttc	tgg	gaa	agc	atg	1996
	Pro	Arg	Leu	Glu	Thr	Cys	Tyr	Glu	Glu	Phe	Cys	Phe	Trp	Glu	Ser	Met	
			605					610					615				

tta tgg cac tgg ggt gaa aca aac tgg aca gat aat gcc acc tct tca 2044
 Leu Trp His Trp Gly Glu Thr Asn Trp Thr Asp Asn Ala Thr Ser Ser
 620 625 630

5 cca aca cot ccc atg gtc aat act gct tca ctt tct tct ttg 2086
 Pro Thr Pro Pro Met Val Asn Thr Ala Ser Leu Ser Ser Leu
 635 640 645

10 taactaggag gttctgttac tatacctcgc tagtctgtaa gtaatacaga gtcacaactt 2146
 gaatgtcaac cagtgccatt agtatacatg gtgtcaacac tttccccacc ettgaaaaaa 2206
 aaaaaaaaaa aaaaaaaaaa aaaa 2230

15 <210> 4
 <211> 648
 <212> PRT
 20 <213> Solanum tuberosum

<400> 4
 Met Arg Gly Ser Leu Ala Gly Gly Pro Pro Ser Pro Ile Glu Pro Arg
 1 5 10 15

25 Gln Arg Leu Ser Val Phe Thr Glu Glu Thr Ser Lys Arg Arg Phe Leu
 20 25 30

30 Arg Ser Lys Val Phe Arg Asp Gly Glu Arg Ala Leu His Ser Pro Thr
 35 40 45

Lys Asn Arg Asn Phe Thr Cys Lys Phe Pro Thr Val Lys Leu Ile Leu
 50 55 60

35 Gly Val Ile Ala Leu Val Ala Ile Trp Ser Leu Trp His Ser Pro Ala
 65 70 75 80

Ile Tyr Asn Thr Glu Tyr Ile Ser Ser Ser Gly Ser Arg Ala Ala Leu
 85 90 95

40 Met His Arg Glu Leu Ser Gly His Ser Ser Ala Asp Gln Arg Tyr Thr
 100 105 110

45 Ser Leu Leu Asp Ile Asp Trp Asp Gln Ile Ser Gln Val Ile Glu Lys
 115 120 125

Leu Ala Asp Arg His Glu Tyr Gln Gly Val Gly Ile Leu Asn Phe Asn
 130 135 140

50 Asp Ser Glu Ile Asp Gln Leu Lys Glu Leu Leu Pro Asp Ala Glu His
 145 150 155 160

Val Ile Leu Asn Leu Asp His Val Pro Asn Asn Ile Thr Trp Glu Thr
 165 170 175

55 Ile Tyr Pro Glu Trp Ile Asp Glu Glu Glu Glu Phe Glu Val Pro Thr
 180 185 190

60 Cys Pro Ser Leu Pro Lys Ile Gln Phe Pro Gly Lys Pro Arg Ile Asp
 195 200 205

Leu Ile Val Val Lys Leu Pro Cys Lys Lys Ser Lys Asp Trp Tyr Arg

210 215 220
 Asp Val Ala Arg Phe His Leu Gln Leu Ala Ala Ala Arg Leu Ala Ala
 225 230 235 240
 5 Ser Asn Lys Gly Tyr His Pro Ile His Val Leu Leu Val Thr Glu His
 245 250 255
 10 Phe Pro Thr Pro Asn Leu Phe Thr Cys Lys Glu Leu Val Val Arg Glu
 260 265 270
 Gly Asn Ala Trp Leu Tyr Glu Pro Asn Leu Asn Thr Leu Arg Glu Lys
 275 280 285
 15 Leu His Leu Pro Val Gly Ser Cys Glu Leu Ala Val Pro Leu Lys Ala
 290 295 300
 Lys Ala Asn Trp His Ser Gly Asn Val Arg Arg Glu Ala Tyr Ala Thr
 305 310 315 320
 20 Ile Leu His Ser Ala Asn Phe Tyr Val Cys Gly Ala Ile Ala Ala Ala
 325 330 335
 Gln Ser Ile Arg Leu Ala Gly Ser Thr Arg Asp Leu Val Ile Leu Val
 340 345 350
 25 Asp Glu Thr Ile Ser Asp Tyr His Arg Gly Gly Leu Glu Ala Ala Gly
 355 360 365
 30 Trp Lys Ile His Thr Ile Lys Arg Ile Arg Asn Pro Lys Ala Glu Gln
 370 375 380
 Asp Ala Tyr Asn Glu Trp Asn Tyr Ser Lys Phe Arg Leu Trp Gln Leu
 385 390 395 400
 35 Thr Asp Tyr Asp Lys Ile Ile Phe Ile Asp Ala Asp Leu Leu Ile Leu
 405 410 415
 Arg Asn Ile Asp Phe Leu Phe Glu Met Pro Glu Ile Thr Ala Ile Gly
 420 425 430
 Asn Asn Ala Thr Leu Phe Asn Ser Gly Val Met Val Val Glu Pro Ser
 435 440 445
 45 Asn Cys Thr Phe Gln Leu Leu Met Asp His Ile Asn Glu Ile Glu Ser
 450 455 460
 Tyr Asn Gly Gly Asp Gln Gly Tyr Leu Asn Glu Ile Phe Thr Trp Trp
 465 470 475 480
 50 His Arg Ile Pro Lys His Met Asn Phe Leu Lys His Tyr Trp Glu Gly
 485 490 495
 Asp Glu Glu Glu Lys Lys Gln Met Lys Thr Arg Leu Phe Gly Ala Asp
 500 505 510
 Pro Pro Val Leu Tyr Val Leu His Tyr Leu Gly Leu Lys Pro Trp Leu
 515 520 525
 60 Cys Phe Arg Asp Tyr Asp Cys Asn Trp Asn Val Gly Lys Leu Gln Glu
 530 535 540

10

	Phe	Ala	Ser	Asp	Val	Ala	His	Arg	Thr	Trp	Trp	Lys	Val	His	Asp	Ala	
	545					550					555					560	
5	Met	Pro	Asp	Asn	Leu	His	Lys	Tyr	Cys	Leu	Leu	Arg	Ser	Lys	Gln	Lys	
					565					570					575		
	Ala	Ala	Leu	Glu	Trp	Asp	Arg	Arg	Glu	Ala	Glu	Lys	Ala	Asn	Phe	Ser	
				580					585					590			
10	Asp	Gly	His	Trp	Lys	Ile	Lys	Ile	Lys	Asp	Pro	Arg	Leu	Glu	Thr	Cys	
			595					600					605				
	Tyr	Glu	Glu	Phe	Cys	Phe	Trp	Glu	Ser	Met	Leu	Trp	His	Trp	Gly	Glu	
	610						615					620					
15	Thr	Asn	Trp	Thr	Asp	Asn	Ala	Thr	Ser	Ser	Pro	Thr	Pro	Pro	Met	Val	
	625					630					635				640		
20	Asn	Thr	Ala	Ser	Leu	Ser	Ser	Leu									
					645												
25	<210> 5																
	<211> 1485																
	<212> DNA																
	<213> Arabidopsis thaliana																
30	<220>																
	<221> CDS																
	<222> (1) .. (1485)																
35	<400> 5																
	atg	gat	tta	caa	aga	act	ttg	atg	ttc	tct	tgt	tgg	gtt	ctg	tct	ctt	48
	Met	Asp	Leu	Gln	Arg	Thr	Leu	Met	Phe	Ser	Cys	Trp	Val	Leu	Ser	Leu	
	1					5				10				15			
40	ttg	atc	atc	aaa	acg	aca	gcg	tat	aac	gag	aaa	cag	ctg	ttc	cag	ccg	96
	Leu	Ile	Ile	Lys	Thr	Thr	Ala	Tyr	Asn	Glu	Lys	Gln	Leu	Phe	Gln	Pro	
				20					25					30			
45	ctt	gaa	acg	gaa	aac	gca	aac	gcg	atg	acc	gcg	gtt	atg	gag	cga	gga	144
	Leu	Glu	Thr	Glu	Asn	Ala	Asn	Ala	Met	Thr	Ala	Val	Met	Glu	Arg	Gly	
				35				40					45				
50	tta	aag	acg	cag	cgg	cgg	ccg	gag	cac	aag	aac	gct	tat	gcg	acg	atg	192
	Leu	Lys	Thr	Gln	Arg	Arg	Pro	Glu	His	Lys	Asn	Ala	Tyr	Ala	Thr	Met	
		50					55					60					
55	atg	tac	atg	gga	aca	cca	aga	gac	tac	gag	ttc	tac	gtt	gcg	aca	cgt	240
	Met	Tyr	Met	Gly	Thr	Pro	Arg	Asp	Tyr	Glu	Phe	Tyr	Val	Ala	Thr	Arg	
	65					70					75					80	
60	gtc	ttg	atc	aga	tcg	ctt	aag	agt	ctc	cac	gtg	gac	gct	gat	atc	gtc	288
	Val	Leu	Ile	Arg	Ser	Leu	Lys	Ser	Leu	His	Val	Asp	Ala	Asp	Ile	Val	
					85					90					95		
65	gtt	ata	gcc	tcc	ctc	gac	gtt	cct	atc	aac	tgg	att	cac	gct	ctg	gaa	336
	Val	Ile	Ala	Ser	Leu	Asp	Val	Pro	Ile	Asn	Trp	Ile	His	Ala	Leu	Glu	
				100					105					110			
70	gaa	gaa	gat	gga	gct	aaa	gta	gtg	aga	gta	gag	aat	ctt	gag	aat	cca	384

11

	Glu Glu Asp Gly Ala Lys Val Val Arg Val Glu Asn Leu Glu Asn Pro	
	115 120 125	
5	tac aag aaa caa acc aac ttc gac aac aga ttc aag ctt agt cta aac Tyr Lys Lys Gln Thr Asn Phe Asp Asn Arg Phe Lys Leu Ser Leu Asn	432
	130 135 140	
10	aag ctc tac gct tgg tct ctc tct gat tat gac cgt gtt gta atg ctt Lys Leu Tyr Ala Trp Ser Leu Ser Asp Tyr Asp Arg Val Val Met Leu	480
	145 150 155 160	
15	gat gtc gac aat ctc ttt ctc aag aac acc gac gag ctc ttc cag tgt Asp Val Asp Asn Leu Phe Leu Lys Asn Thr Asp Glu Leu Phe Gln Cys	528
	165 170 175	
	ggc caa ttt tgt gct gtc ttc atc aac cct tgc atc ttc cac act ggt Gly Gln Phe Cys Ala Val Phe Ile Asn Pro Cys Ile Phe His Thr Gly	576
	180 185 190	
20	ctc ttt gtg ttg cag cca tca atg gag gtc ttt aga gac atg ctt cat Leu Phe Val Leu Gln Pro Ser Met Glu Val Phe Arg Asp Met Leu His	624
	195 200 205	
25	gag ctt gaa gta aag aga gat aac cct gat gga gct gat caa ggc ttt Glu Leu Glu Val Lys Arg Asp Asn Pro Asp Gly Ala Asp Gln Gly Phe	672
	210 215 220	
30	ctt gtc agc tac ttc tct gat tta ctc aat cag cct ctc ttt cgt cct Leu Val Ser Tyr Phe Ser Asp Leu Leu Asn Gln Pro Leu Phe Arg Pro	720
	225 230 235 240	
35	cct ccc gat aac cgc acc gcg ctt aag gga cat ttt agg ctt cct ttg Pro Pro Asp Asn Arg Thr Ala Leu Lys Gly His Phe Arg Leu Pro Leu	768
	245 250 255	
	gga tat caa atg gac gca tct tat tac tac ctt aag ctc aga tgg aac Gly Tyr Gln Met Asp Ala Ser Tyr Tyr Tyr Leu Lys Leu Arg Trp Asn	816
	260 265 270	
40	gta cca tgt gga cca aac agt gtg ata acg ttc cca gga gca gta tgg Val Pro Cys Gly Pro Asn Ser Val Ile Thr Phe Pro Gly Ala Val Trp	864
	275 280 285	
45	tta aag cca tgg tat tgg tgg tca tgg cct gtt ctt cct tta ggc ctt Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro Leu Gly Leu	912
	290 295 300	
50	tca tgg cac cac caa cgc cgc tac acg att agt tat tca gca gag atg Ser Trp His His Gln Arg Arg Tyr Thr Ile Ser Tyr Ser Ala Glu Met	960
	305 310 315 320	
55	cct tgg gtc cta acc caa gca gtg ttc tac cta gga att ata cta gtc Pro Trp Val Leu Thr Gln Ala Val Phe Tyr Leu Gly Ile Ile Leu Val	1008
	325 330 335	
	aca cgt cta gcg cgt ccc aac atg acc aag cta tgt tac cga cgt tct Thr Arg Leu Ala Arg Pro Asn Met Thr Lys Leu Cys Tyr Arg Arg Ser	1056
	340 345 350	
60	gat aag aat cta agc atg atc cag aca gct ttc aag ttt gtt gca ctc Asp Lys Asn Leu Ser Met Ile Gln Thr Ala Phe Lys Phe Val Ala Leu	1104
	355 360 365	

12

5	ctc ttt atc ctc tca gcc tac att ata cca ttc ttc atc atc cca cag	1152
	Leu Phe Ile Leu Ser Ala Tyr Ile Ile Pro Phe Phe Ile Ile Pro Gln	
	370 375 380	
	acg atc cac ccg ctc att ggt tgg tot ctc tac tta acc ggc tcc ttt	1200
	Thr Ile His Pro Leu Ile Gly Trp Ser Leu Tyr Leu Thr Gly Ser Phe	
10	gct ctc tct acc ata ccc atc aac gcc ttc ttg ctt ccc att ctc cct	1248
	Ala Leu Ser Thr Ile Pro Ile Asn Ala Phe Leu Leu Pro Ile Leu Pro	
	405 410 415	
	gtc ata aca ccg tgg ctt ggc att ttc ggg aca ctc ctc gtg atg gct	1296
	Val Ile Thr Pro Trp Leu Gly Ile Phe Gly Thr Leu Leu Val Met Ala	
20	ttt cct tct tat cct gat ggc gtt gtt aga gca ttg tcg gtt ttc ggg	1344
	Phe Pro Ser Tyr Pro Asp Gly Val Val Arg Ala Leu Ser Val Phe Gly	
	435 440 445	
	tat gca ttt tgt tgt gca ccg ttt cta tgg gtc tcc ttt gtg aag atc	1392
	Tyr Ala Phe Cys Cys Ala Pro Phe Leu Trp Val Ser Phe Val Lys Ile	
25	aca tcg cat ctt cag att atg att gac aaa gag gtt ttg ttt ccg cgg	1440
	Thr Ser His Leu Gln Ile Met Ile Asp Lys Glu Val Leu Phe Pro Arg	
	465 470 475 480	
	ttg ggt gaa tcc gga gtc act tct ggt ctc agc aaa ttg tac tga	1485
	Leu Gly Glu Ser Gly Val Thr Ser Gly Leu Ser Lys Leu Tyr	
30	485 490 495	
35	<210> 6	
	<211> 494	
	<212> PRT	
	<213> Arabidopsis thaliana	
40	<400> 6	
	Met Asp Leu Gln Arg Thr Leu Met Phe Ser Cys Trp Val Leu Ser Leu	
	1 5 10 15	
	Leu Ile Ile Lys Thr Thr Ala Tyr Asn Glu Lys Gln Leu Phe Gln Pro	
	20 25 30	
45	Leu Glu Thr Glu Asn Ala Asn Ala Met Thr Ala Val Met Glu Arg Gly	
	35 40 45	
	Leu Lys Thr Gln Arg Arg Pro Glu His Lys Asn Ala Tyr Ala Thr Met	
	50 55 60	
	Met Tyr Met Gly Thr Pro Arg Asp Tyr Glu Phe Tyr Val Ala Thr Arg	
50	65 70 75 80	
	Val Leu Ile Arg Ser Leu Lys Ser Leu His Val Asp Ala Asp Ile Val	
	85 90 95	
	Val Ile Ala Ser Leu Asp Val Pro Ile Asn Trp Ile His Ala Leu Glu	
	100 105 110	
55	Glu Glu Asp Gly Ala Lys Val Val Arg Val Glu Asn Leu Glu Asn Pro	
	115 120 125	
	Tyr Lys Lys Gln Thr Asn Phe Asp Asn Arg Phe Lys Leu Ser Leu Asn	
	130 135 140	
	Lys Leu Tyr Ala Trp Ser Leu Ser Asp Tyr Asp Arg Val Val Met Leu	
60	145 150 155 160	
	Asp Val Asp Asn Leu Phe Leu Lys Asn Thr Asp Glu Leu Phe Gln Cys	
	165 170 175	

13

Gly Gln Phe Cys Ala Val Phe Ile Asn Pro Cys Ile Phe His Thr Gly
 180 185 190
 Leu Phe Val Leu Gln Pro Ser Met Glu Val Phe Arg Asp Met Leu His
 195 200 205
 5 Glu Leu Glu Val Lys Arg Asp Asn Pro Asp Gly Ala Asp Gln Gly Phe
 210 215 220
 Leu Val Ser Tyr Phe Ser Asp Leu Leu Asn Gln Pro Leu Phe Arg Pro
 225 230 235 240
 10 Pro Pro Asp Asn Arg Thr Ala Leu Lys Gly His Phe Arg Leu Pro Leu
 245 250 255
 Gly Tyr Gln Met Asp Ala Ser Tyr Tyr Tyr Leu Lys Leu Arg Trp Asn
 260 265 270
 Val Pro Cys Gly Pro Asn Ser Val Ile Thr Phe Pro Gly Ala Val Trp
 275 280 285
 15 Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro Leu Gly Leu
 290 295 300
 Ser Trp His His Gln Arg Arg Tyr Thr Ile Ser Tyr Ser Ala Glu Met
 305 310 315 320
 20 Pro Trp Val Leu Thr Gln Ala Val Phe Tyr Leu Gly Ile Ile Leu Val
 325 330 335
 Thr Arg Leu Ala Arg Pro Asn Met Thr Lys Leu Cys Tyr Arg Arg Ser
 340 345 350
 Asp Lys Asn Leu Ser Met Ile Gln Thr Ala Phe Lys Phe Val Ala Leu
 355 360 365
 25 Leu Phe Ile Leu Ser Ala Tyr Ile Ile Pro Phe Phe Ile Ile Pro Gln
 370 375 380
 Thr Ile His Pro Leu Ile Gly Trp Ser Leu Tyr Leu Thr Gly Ser Phe
 385 390 395 400
 30 Ala Leu Ser Thr Ile Pro Ile Asn Ala Phe Leu Leu Pro Ile Leu Pro
 405 410 415
 Val Ile Thr Pro Trp Leu Gly Ile Phe Gly Thr Leu Leu Val Met Ala
 420 425 430
 Phe Pro Ser Tyr Pro Asp Gly Val Val Arg Ala Leu Ser Val Phe Gly
 435 440 445
 35 Tyr Ala Phe Cys Cys Ala Pro Phe Leu Trp Val Ser Phe Val Lys Ile
 450 455 460
 Thr Ser His Leu Gln Ile Met Ile Asp Lys Glu Val Leu Phe Pro Arg
 465 470 475 480
 40 Leu Gly Glu Ser Gly Val Thr Ser Gly Leu Ser Lys Leu Tyr
 485 490

45 <210> 7
 <211> 1494
 <212> DNA
 <213> Arabidopsis thaliana

50 <220>
 <221> CDS
 <222> (1)..(1494)

55 <400> 7
 atg gat ttg cag aga ggc ttt gtg ttc ttg tot ttg gtt cta tct ttt 48
 Met Asp Leu Gln Arg Gly Phe Val Phe Leu Ser Leu Val Leu Ser Phe
 1 5 10 15

60 atg ata atc gaa acg acg gcg tat cga gag aga cag ctg ctg ctg ctg 96
 Met Ile Ile Glu Thr Thr Ala Tyr Arg Glu Arg Gln Leu Leu Leu
 20 25 30

caa cca ccg caa gaa acg gcg ata gat acc gca aac gcg gtg gtg acg 144

14

	Gln	Pro	Pro	Gln	Glu	Thr	Ala	Ile	Asp	Thr	Ala	Asn	Ala	Val	Val	Thr	
		35						40					45				
5	gtt	caa	gat	cga	ggt	ttg	aag	acg	cgg	cga	ccg	gag	cat	aag	aac	gca	192
	Val	Gln	Asp	Arg	Gly	Leu	Lys	Thr	Arg	Arg	Pro	Glu	His	Lys	Asn	Ala	
		50				55					60						
10	tac	gca	acg	atg	atg	tac	atg	ggg	acg	cca	aga	gac	tac	gag	ttc	tac	240
	Tyr	Ala	Thr	Met	Met	Tyr	Met	Gly	Thr	Pro	Arg	Asp	Tyr	Glu	Phe	Tyr	
		65				70				75					80		
15	gtt	gcg	aca	cgt	gtt	ttg	atc	aga	tcg	ttg	aga	agt	ctt	cac	gtg	gaa	288
	Val	Ala	Thr	Arg	Val	Leu	Ile	Arg	Ser	Leu	Arg	Ser	Leu	His	Val	Glu	
				85					90						95		
	gct	gat	ctc	gtc	gtc	atc	gct	tct	ctc	gac	gtt	cct	ctc	cga	tgg	gtt	336
	Ala	Asp	Leu	Val	Val	Ile	Ala	Ser	Leu	Asp	Val	Pro	Leu	Arg	Trp	Val	
				100					105					110			
20	caa	acc	ttg	gaa	gag	gaa	gat	gga	gct	aaa	gtg	gtg	aga	gtt	gaa	aat	384
	Gln	Thr	Leu	Glu	Glu	Glu	Asp	Gly	Ala	Lys	Val	Val	Arg	Val	Glu	Asn	
			115					120					125				
25	gtg	gat	aat	cca	tac	agg	aga	cag	acc	aac	ttc	aac	agt	aga	ttc	aag	432
	Val	Asp	Asn	Pro	Tyr	Arg	Arg	Gln	Thr	Asn	Phe	Asn	Ser	Arg	Phe	Lys	
		130					135					140					
30	ctt	act	cta	aac	aag	ctc	tac	gct	tgg	gct	ttg	tct	gat	tac	gac	cgt	480
	Leu	Thr	Leu	Asn	Lys	Leu	Tyr	Ala	Trp	Ala	Leu	Ser	Asp	Tyr	Asp	Arg	
		145				150					155					160	
35	gtg	gtc	atg	cta	gat	gcc	gat	aac	ctc	ttt	ctt	aag	aaa	gcc	gac	gag	528
	Val	Val	Met	Leu	Asp	Ala	Asp	Asn	Leu	Phe	Leu	Lys	Lys	Ala	Asp	Glu	
				165					170						175		
	ttg	ttc	cag	tgt	ggg	cgc	ttc	tgt	gcg	gtc	ttc	att	aac	cct	tgt	atc	576
	Leu	Phe	Gln	Cys	Gly	Arg	Phe	Cys	Ala	Val	Phe	Ile	Asn	Pro	Cys	Ile	
			180						185					190			
40	ttc	cac	act	ggt	ctc	ttc	gtg	ttg	cag	cca	tca	gtg	gaa	gtg	ttc	aag	624
	Phe	His	Thr	Gly	Leu	Phe	Val	Leu	Gln	Pro	Ser	Val	Glu	Val	Phe	Lys	
			195					200					205				
45	gac	atg	ctc	cat	gag	cta	caa	gtt	gga	aga	aag	aat	cct	gat	gga	gct	672
	Asp	Met	Leu	His	Glu	Leu	Gln	Val	Gly	Arg	Lys	Asn	Pro	Asp	Gly	Ala	
		210					215					220					
50	gat	caa	ggt	ttt	ctt	gtc	agt	tac	ttc	tct	gat	ctt	ctt	gac	caa	cct	720
	Asp	Gln	Gly	Phe	Leu	Val	Ser	Tyr	Phe	Ser	Asp	Leu	Leu	Asp	Gln	Pro	
		225				230					235					240	
55	ctc	ttt	agt	cct	ccg	agt	aac	gga	tct	gta	ctt	aat	ggt	cac	ttg	aga	768
	Leu	Phe	Ser	Pro	Pro	Ser	Asn	Gly	Ser	Val	Leu	Asn	Gly	His	Leu	Arg	
				245						250					255		
	ctt	ccc	tta	ggc	tac	caa	atg	gac	gct	tct	tat	ttc	tat	ctt	aag	cta	816
	Leu	Pro	Leu	Gly	Tyr	Gln	Met	Asp	Ala	Ser	Tyr	Phe	Tyr	Leu	Lys	Leu	
				260					265						270		
60	aga	tgg	aac	ata	ccc	tgt	gga	cca	aac	agt	gtg	att	aca	ttc	ccg	gga	864
	Arg	Trp	Asn	Ile	Pro	Cys	Gly	Pro	Asn	Ser	Val	Ile	Thr	Phe	Pro	Gly	
			275					280						285			

	get gtt tgg tta aag cca tgg tac tgg tgg tca tgg cct gtt ctt cca	912
	Ala Val Trp Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro	
	290 295 300	
5	cta ggt ttc tca tgg cac gag cag cgt cgc gcc act ata ggg tac tca	960
	Leu Gly Phe Ser Trp His Glu Gln Arg Arg Ala Thr Ile Gly Tyr Ser	
	305 310 315 320	
10	gcc gaa atg cct ttg gtt ata atc caa gca atg ttt tac ctt gga atc	1008
	Ala Glu Met Pro Leu Val Ile Ile Gln Ala Met Phe Tyr Leu Gly Ile	
	325 330 335	
15	ata gtg gtt aca cga cta gct cgt cct aac ata acc aag cta tgt tat	1056
	Ile Val Val Thr Arg Leu Ala Arg Pro Asn Ile Thr Lys Leu Cys Tyr	
	340 345 350	
20	cgc cgc tct gac cgc aac tta aca acg atc caa gct ggt ttt aag ttg	1104
	Arg Arg Ser Asp Arg Asn Leu Thr Thr Ile Gln Ala Gly Phe Lys Leu	
	355 360 365	
25	atc gcg ctt ctc tct gta gtt gca gcc tac atc ttc cca ttc ttc acc	1152
	Ile Ala Leu Leu Ser Val Val Ala Ala Tyr Ile Phe Pro Phe Phe Thr	
	370 375 380	
30	atc cct cac act atc cac cca ctc atc ggc tgg tgg ctc tac ttg atg	1200
	Ile Pro His Thr Ile His Pro Leu Ile Gly Trp Ser Leu Tyr Leu Met	
	385 390 395 400	
35	gct tct ttc gct ctc tct tcc att tca atc aac act ctc ctc ctc cca	1248
	Ala Ser Phe Ala Leu Ser Ser Ile Ser Ile Asn Thr Leu Leu Leu Pro	
	405 410 415	
40	acg ctc cct gtt ctc act cca tgg cta gga att ctc ggc act ctc ctt	1296
	Thr Leu Pro Val Leu Thr Pro Trp Leu Gly Ile Leu Gly Thr Leu Leu	
	420 425 430	
45	gtc atg gcc ttc cct tgg tac cct gat gga gtg gtc aga gcc ttg tca	1344
	Val Met Ala Phe Pro Trp Tyr Pro Asp Gly Val Val Arg Ala Leu Ser	
	435 440 445	
50	gtt ttc gca tac gca ttt tgt tgc gca ccc ttt gtg tgg gtt tca ttc	1392
	Val Phe Ala Tyr Ala Phe Cys Cys Ala Pro Phe Val Trp Val Ser Phe	
	450 455 460	
55	cgc aaa atc aca tgg cac ctc cag gtt ttg att gag aaa gag gtg ttg	1440
	Arg Lys Ile Thr Ser His Leu Gln Val Leu Ile Glu Lys Glu Val Leu	
	465 470 475 480	
60	ttc ccg cga ttg gga gac tca ggg gtc act tca ggc ttc agc aaa ttg	1488
	Phe Pro Arg Leu Gly Asp Ser Gly Val Thr Ser Gly Phe Ser Lys Leu	
	485 490 495	
65	tat tag	
	Tyr	1494
	<210> 8	
	<211> 497	
	<212> PRT	
	<213> Arabidopsis thaliana	

<400> 8
 Met Asp Leu Gln Arg Gly Phe Val Phe Leu Ser Leu Val Leu Ser Phe
 1 5 10 15
 5 Met Ile Ile Glu Thr Thr Ala Tyr Arg Glu Arg Gln Leu Leu Leu
 20 25 30
 Gln Pro Pro Gln Glu Thr Ala Ile Asp Thr Ala Asn Ala Val Val Thr
 35 40 45
 Val Gln Asp Arg Gly Leu Lys Thr Arg Arg Pro Glu His Lys Asn Ala
 50 55 60
 10 Tyr Ala Thr Met Met Tyr Met Gly Thr Pro Arg Asp Tyr Glu Phe Tyr
 65 70 75 80
 Val Ala Thr Arg Val Leu Ile Arg Ser Leu Arg Ser Leu His Val Glu
 85 90 95
 15 Ala Asp Leu Val Val Ile Ala Ser Leu Asp Val Pro Leu Arg Trp Val
 100 105 110
 Gln Thr Leu Glu Glu Glu Asp Gly Ala Lys Val Val Arg Val Glu Asn
 115 120 125
 Val Asp Asn Pro Tyr Arg Arg Gln Thr Asn Phe Asn Ser Arg Phe Lys
 130 135 140
 20 Leu Thr Leu Asn Lys Leu Tyr Ala Trp Ala Leu Ser Asp Tyr Asp Arg
 145 150 155 160
 Val Val Met Leu Asp Ala Asp Asn Leu Phe Leu Lys Lys Ala Asp Glu
 165 170 175
 25 Leu Phe Gln Cys Gly Arg Phe Cys Ala Val Phe Ile Asn Pro Cys Ile
 180 185 190
 Phe His Thr Gly Leu Phe Val Leu Gln Pro Ser Val Glu Val Phe Lys
 195 200 205
 Asp Met Leu His Glu Leu Gln Val Gly Arg Lys Asn Pro Asp Gly Ala
 210 215 220
 30 Asp Gln Gly Phe Leu Val Ser Tyr Phe Ser Asp Leu Leu Asp Gln Pro
 225 230 235 240
 Leu Phe Ser Pro Pro Ser Asn Gly Ser Val Leu Asn Gly His Leu Arg
 245 250 255
 35 Leu Pro Leu Gly Tyr Gln Met Asp Ala Ser Tyr Phe Tyr Leu Lys Leu
 260 265 270
 Arg Trp Asn Ile Pro Cys Gly Pro Asn Ser Val Ile Thr Phe Pro Gly
 275 280 285
 Ala Val Trp Leu Lys Pro Trp Tyr Trp Trp Ser Trp Pro Val Leu Pro
 290 295 300
 40 Leu Gly Phe Ser Trp His Glu Gln Arg Arg Ala Thr Ile Gly Tyr Ser
 305 310 315 320
 Ala Glu Met Pro Leu Val Ile Ile Gln Ala Met Phe Tyr Leu Gly Ile
 325 330 335
 45 Ile Val Val Thr Arg Leu Ala Arg Pro Asn Ile Thr Lys Leu Cys Tyr
 340 345 350
 Arg Arg Ser Asp Arg Asn Leu Thr Thr Ile Gln Ala Gly Phe Lys Leu
 355 360 365
 Ile Ala Leu Leu Ser Val Val Ala Ala Tyr Ile Phe Pro Phe Phe Thr
 370 375 380
 50 Ile Pro His Thr Ile His Pro Leu Ile Gly Trp Ser Leu Tyr Leu Met
 385 390 395 400
 Ala Ser Phe Ala Leu Ser Ser Ile Ser Ile Asn Thr Leu Leu Leu Pro
 405 410 415
 55 Thr Leu Pro Val Leu Thr Pro Trp Leu Gly Ile Leu Gly Thr Leu Leu
 420 425 430
 Val Met Ala Phe Pro Trp Tyr Pro Asp Gly Val Val Arg Ala Leu Ser
 435 440 445
 Val Phe Ala Tyr Ala Phe Cys Cys Ala Pro Phe Val Trp Val Ser Phe
 450 455 460
 60 Arg Lys Ile Thr Ser His Leu Gln Val Leu Ile Glu Lys Glu Val Leu
 465 470 475 480
 Phe Pro Arg Leu Gly Asp Ser Gly Val Thr Ser Gly Phe Ser Lys Leu

17

Tyr

485

490

495

5

<210> 9

<211> 1944

<212> DNA

<213> Arabidopsis thaliana

10

<220>

<221> CDS

<222> (1) .. (1944)

15

<400> 9

atg cat cca gct tca tgt agt ctc tca ctc agg aaa gtc aaa ctt aat 48
 Met His Pro Ala Ser Cys Ser Leu Ser Leu Arg Lys Val Lys Leu Asn
 1 5 10 15

20

att ctt aca gtg aga atg aag ctc tct tct gaa agt ccc atg gcg ccg 96
 Ile Leu Thr Val Arg Met Lys Leu Ser Ser Glu Ser Pro Met Ala Pro
 20 25 30

25

tca agt cag tca agt cat cga ctt tac att tcc agt gag aaa aca aaa 144
 Ser Ser Gln Ser Ser His Arg Leu Tyr Ile Ser Ser Glu Lys Thr Lys
 35 40 45

30

acg aag aga ttc caa aga aac gga tac act ctc gat gtt gaa atg tgt 192
 Thr Lys Arg Phe Gln Arg Asn Gly Tyr Thr Leu Asp Val Glu Met Cys
 50 55 60

35

gtc aac ttc tct agt ctg aaa ctt gtt ttg ttt ctt atg atg ctg gtt 240
 Val Asn Phe Ser Ser Leu Lys Leu Val Leu Phe Leu Met Met Leu Val
 65 70 75 80

gct atg ttc aca ctc tac tgt tct cca ccg ttg caa att cct gaa gat 288
 Ala Met Phe Thr Leu Tyr Cys Ser Pro Pro Leu Gln Ile Pro Glu Asp
 85 90 95

40

cca tca agt ttt gca aac aaa tgg ata cta gaa cct gct gta acc aca 336
 Pro Ser Ser Phe Ala Asn Lys Trp Ile Leu Glu Pro Ala Val Thr Thr
 100 105 110

45

gat cct cgc tat ata gct aca tot gag atc aac tgg aac agt atg tca 384
 Asp Pro Arg Tyr Ile Ala Thr Ser Glu Ile Asn Trp Asn Ser Met Ser
 115 120 125

50

ctt gtt gtt gag cat tac tta tct ggc aga agc gag tat caa gga att 432
 Leu Val Val Glu His Tyr Leu Ser Gly Arg Ser Glu Tyr Gln Gly Ile
 130 135 140

55

ggc ttt cta aat ctc aac gat aac gag att aat cga tgg cag gtg gtc 480
 Gly Phe Leu Asn Leu Asn Asp Asn Glu Ile Asn Arg Trp Gln Val Val
 145 150 155 160

ata aaa tct cac tgt cag cat ata gct ttg cat cta gac cat gct gca 528
 Ile Lys Ser His Cys Gln His Ile Ala Leu His Leu Asp His Ala Ala
 165 170 175

60

agt aac ata act tgg aaa tct tta tac ccg gaa tgg att gac gag gaa 576
 Ser Asn Ile Thr Trp Lys Ser Leu Tyr Pro Glu Trp Ile Asp Glu Glu
 180 185 190

5	gaa aaa ttc aaa gtc ccc act tgt cct tct ctt cct tgg att caa gtt Glu Lys Phe Lys Val Pro Thr Cys Pro Ser Leu Pro Trp Ile Gln Val 195 200 205	624
10	cct gac aag tct cga atc gat ctt atc att gcc aag ctc cca tgt aac Pro Asp Lys Ser Arg Ile Asp Leu Ile Ile Ala Lys Leu Pro Cys Asn 210 215 220	672
15	aag tca gga aaa tgg tca aga gat gtg gct aga ttg cac tta caa ctt Lys Ser Gly Lys Trp Ser Arg Asp Val Ala Arg Leu His Leu Gln Leu 225 230 235 240	720
20	gca gca gct cga gtg gcg gca tct tct gaa ggg ctt cat gat gtt cat Ala Ala Ala Arg Val Ala Ala Ser Ser Glu Gly Leu His Asp Val His 245 250 255	768
25	gtg att ttg gta tca gat tgc ttt cca ata ccg aat ctt ttt acg ggt Val Ile Leu Val Ser Asp Cys Phe Pro Ile Pro Asn Leu Phe Thr Gly 260 265 270	816
30	caa gaa ctt gtt gcc cgt caa gga aac ata tgg ctg tat aag cct aaa Gln Glu Leu Val Ala Arg Gln Gly Asn Ile Trp Leu Tyr Lys Pro Lys 275 280 285	864
35	ctt cac cag tta aga caa aag tta caa ctt cct gtt ggt tcc tgt gaa Leu His Gln Leu Arg Gln Lys Leu Gln Leu Pro Val Gly Ser Cys Glu 290 295 300	912
40	ctt tct gtt cct ctt caa gct aaa gat aat ttc tac tcg gca aat gcc Leu Ser Val Pro Leu Gln Ala Lys Asp Asn Phe Tyr Ser Ala Asn Ala 305 310 315 320	960
45	aag aaa gaa gog tac gcg acg atc ttg cac tca gat gat gct ttt gtc Lys Lys Glu Ala Tyr Ala Thr Ile Leu His Ser Asp Asp Ala Phe Val 325 330 335	1008
50	tgt gga gcc att gca gta gca cag agc att cga atg tca ggc tct act Cys Gly Ala Ile Ala Val Ala Gln Ser Ile Arg Met Ser Gly Ser Thr 340 345 350	1056
55	cgc aat ttg gta ata cta gtc gat gat tcg atc agt gaa tac cat aga Arg Asn Leu Val Ile Leu Val Asp Asp Ser Ile Ser Glu Tyr His Arg 355 360 365	1104
60	agt ggc ttg gaa tca gct gga tgg aag att cac aca ttt caa aga atc Ser Gly Leu Glu Ser Ala Gly Trp Lys Ile His Thr Phe Gln Arg Ile 370 375 380	1152
65	aga aac ccg aaa gct gaa gca aat gca tat aac caa tgg aac tac agc Arg Asn Pro Lys Ala Glu Ala Asn Ala Tyr Asn Gln Trp Asn Tyr Ser 385 390 395 400	1200
70	aaa ttc cgt ctt tgg gaa ttg aca gaa tac aac aag atc atc ttc att Lys Phe Arg Leu Trp Glu Leu Thr Glu Tyr Asn Lys Ile Ile Phe Ile 405 410 415	1248
75	gat gca gac atg ctt atc ctc aga aac atg gat ttc ctc ttc gag tac Asp Ala Asp Met Leu Ile Leu Arg Asn Met Asp Phe Leu Phe Glu Tyr 420 425 430	1296

19

	ccc gaa atc tcc aca act gga aac gac ggt acg ctc ttc aac tcc ggt Pro Glu Ile Ser Thr Thr Gly Asn Asp Gly Thr Leu Phe Asn Ser Gly	1344
	435 440 445	
5	cta atg gtg att gaa cca tca aat tca aca ttc cag tta cta atg gat Leu Met Val Ile Glu Pro Ser Asn Ser Thr Phe Gln Leu Leu Met Asp	1392
	450 455 460	
10	cac atc aac gat atc aat tcc tac aat gga gga gac caa ggt tac ctt His Ile Asn Asp Ile Asn Ser Tyr Asn Gly Gly Asp Gln Gly Tyr Leu	1440
	465 470 475 480	
15	aac gag ata ttc aca tgg tgg cat cgg att cca aaa cac atg aat ttc Asn Glu Ile Phe Thr Trp Trp His Arg Ile Pro Lys His Met Asn Phe	1488
	485 490 495	
20	tgg aag cat ttc tgg gaa gga gac aca cct aag cac agg aaa tct aag Leu Lys His Phe Thr Glu Gly Asp Thr Pro Lys His Arg Lys Ser Lys	1536
	500 505 510	
	acg aga cta ttt gga gct gat cct ccg ata ctc tac gtt ctt cat tac Thr Arg Leu Phe Gly Ala Asp Pro Pro Ile Leu Tyr Val Leu His Tyr	1584
	515 520 525	
25	cta ggt tac aac aaa cca tgg gta tgc ttc aga gac tac gat tgc aat Leu Gly Tyr Asn Lys Pro Trp Val Cys Phe Arg Asp Tyr Asp Cys Asn	1632
	530 535 540	
30	tgg aat gtc gtt gga tac cat caa ttc gcg agc gat gaa gca cac aaa Trp Asn Val Val Gly Tyr His Gln Phe Ala Ser Asp Glu Ala His Lys	1680
	545 550 555 560	
35	act tgg tgg aga gtg cac gac gcg atg cct aag aaa ttg cag agg ttt Thr Trp Trp Arg Val His Asp Ala Met Pro Lys Lys Leu Gln Arg Phe	1728
	565 570 575	
40	tgt cta ctg agt tcc aaa caa aag gcg caa ctt gag tgg gat cgg aga Cys Leu Leu Ser Ser Lys Gln Lys Ala Gln Leu Glu Trp Asp Arg Arg	1776
	580 585 590	
	caa gct gag aaa gcg aat tac aga gac gga cat tgg agg att aag atc Gln Ala Glu Lys Ala Asn Tyr Arg Asp Gly His Trp Arg Ile Lys Ile	1824
	595 600 605	
45	aaa gat aag aga ctt acg act tgt ttt gaa gat ttc tgt ttc tgg gag Lys Asp Lys Arg Leu Thr Thr Cys Phe Glu Asp Phe Cys Phe Trp Glu	1872
	610 615 620	
50	agt atg ctt tgg cat tgg ggt gag act cag acc aac tcc acc gtc gct Ser Met Leu Trp His Trp Gly Glu Thr Gln Thr Asn Ser Thr Val Ala	1920
	625 630 635 640	
55	gct gat tct tcc tcc acc gcg taa Ala Asp Ser Ser Ser Thr Ala	1944
	645	
60	<210> 10 <211> 647 <212> PRT <213> Arabidopsis thaliana	

20

<400> 10

	Met	His	Pro	Ala	Ser	Cys	Ser	Leu	Ser	Leu	Arg	Lys	Val	Lys	Leu	Asn
	1				5					10					15	
5	Ile	Leu	Thr	Val	Arg	Met	Lys	Leu	Ser	Ser	Glu	Ser	Pro	Met	Ala	Pro
				20				25					30			
	Ser	Ser	Gln	Ser	Ser	His	Arg	Leu	Tyr	Ile	Ser	Ser	Glu	Lys	Thr	Lys
			35					40					45			
	Thr	Lys	Arg	Phe	Gln	Arg	Asn	Gly	Tyr	Thr	Leu	Asp	Val	Glu	Met	Cys
		50					55					60				
10	Val	Asn	Phe	Ser	Ser	Leu	Lys	Leu	Val	Leu	Phe	Leu	Met	Met	Leu	Val
	65					70					75				80	
	Ala	Met	Phe	Thr	Leu	Tyr	Cys	Ser	Pro	Pro	Leu	Gln	Ile	Pro	Glu	Asp
					85					90					95	
15	Pro	Ser	Ser	Phe	Ala	Asn	Lys	Trp	Ile	Leu	Glu	Pro	Ala	Val	Thr	Thr
				100					105					110		
	Asp	Pro	Arg	Tyr	Ile	Ala	Thr	Ser	Glu	Ile	Asn	Trp	Asn	Ser	Met	Ser
			115					120					125			
	Leu	Val	Val	Glu	His	Tyr	Leu	Ser	Gly	Arg	Ser	Glu	Tyr	Gln	Gly	Ile
		130					135					140				
20	Gly	Phe	Leu	Asn	Leu	Asn	Asp	Asn	Glu	Ile	Asn	Arg	Trp	Gln	Val	Val
	145					150						155			160	
	Ile	Lys	Ser	His	Cys	Gln	His	Ile	Ala	Leu	His	Leu	Asp	His	Ala	Ala
				165					170					175		
25	Ser	Asn	Ile	Thr	Trp	Lys	Ser	Leu	Tyr	Pro	Glu	Trp	Ile	Asp	Glu	Glu
			180						185					190		
	Glu	Lys	Phe	Lys	Val	Pro	Thr	Cys	Pro	Ser	Leu	Pro	Trp	Ile	Gln	Val
			195					200					205			
	Pro	Asp	Lys	Ser	Arg	Ile	Asp	Leu	Ile	Ile	Ala	Lys	Leu	Pro	Cys	Asn
		210					215					220				
30	Lys	Ser	Gly	Lys	Trp	Ser	Arg	Asp	Val	Ala	Arg	Leu	His	Leu	Gln	Leu
	225					230					235				240	
	Ala	Ala	Ala	Arg	Val	Ala	Ala	Ser	Ser	Glu	Gly	Leu	His	Asp	Val	His
				245						250				255		
35	Val	Ile	Leu	Val	Ser	Asp	Cys	Phe	Pro	Ile	Pro	Asn	Leu	Phe	Thr	Gly
			260						265					270		
	Gln	Glu	Leu	Val	Ala	Arg	Gln	Gly	Asn	Ile	Trp	Leu	Tyr	Lys	Pro	Lys
			275					280					285			
	Leu	His	Gln	Leu	Arg	Gln	Lys	Leu	Gln	Leu	Pro	Val	Gly	Ser	Cys	Glu
		290					295					300				
40	Leu	Ser	Val	Pro	Leu	Gln	Ala	Lys	Asp	Asn	Phe	Tyr	Ser	Ala	Asn	Ala
	305					310					315				320	
	Lys	Lys	Glu	Ala	Tyr	Ala	Thr	Ile	Leu	His	Ser	Asp	Asp	Ala	Phe	Val
				325						330				335		
45	Cys	Gly	Ala	Ile	Ala	Val	Ala	Gln	Ser	Ile	Arg	Met	Ser	Gly	Ser	Thr
			340						345					350		
	Arg	Asn	Leu	Val	Ile	Leu	Val	Asp	Asp	Ser	Ile	Ser	Glu	Tyr	His	Arg
			355					360					365			
	Ser	Gly	Leu	Glu	Ser	Ala	Gly	Trp	Lys	Ile	His	Thr	Phe	Gln	Arg	Ile
		370					375					380				
50	Arg	Asn	Pro	Lys	Ala	Glu	Ala	Asn	Ala	Tyr	Asn	Gln	Trp	Asn	Tyr	Ser
	385					390					395				400	
	Lys	Phe	Arg	Leu	Trp	Glu	Leu	Thr	Glu	Tyr	Asn	Lys	Ile	Ile	Phe	Ile
				405						410				415		
55	Asp	Ala	Asp	Met	Leu	Ile	Leu	Arg	Asn	Met	Asp	Phe	Leu	Phe	Glu	Tyr
			420						425				430			
	Pro	Glu	Ile	Ser	Thr	Thr	Gly	Asn	Asp	Gly	Thr	Leu	Phe	Asn	Ser	Gly
			435					440				445				
	Leu	Met	Val	Ile	Glu	Pro	Ser	Asn	Ser	Thr	Phe	Gln	Leu	Leu	Met	Asp
		450				455					460					
60	His	Ile	Asn	Asp	Ile	Asn	Ser	Tyr	Asn	Gly	Gly	Asp	Gln	Gly	Tyr	Leu
	465					470					475				480	
	Asn	Glu	Ile	Phe	Thr	Trp	Trp	His	Arg	Ile	Pro	Lys	His	Met	Asn	Phe

21

485 490 495
 Leu Lys His Phe Trp Glu Gly Asp Thr Pro Lys His Arg Lys Ser Lys
 500 505 510
 5 Thr Arg Leu Phe Gly Ala Asp Pro Pro Ile Leu Tyr Val Leu His Tyr
 515 520 525
 5 Leu Gly Tyr Asn Lys Pro Trp Val Cys Phe Arg Asp Tyr Asp Cys Asn
 530 535 540
 Trp Asn Val Val Gly Tyr His Gln Phe Ala Ser Asp Glu Ala His Lys
 545 550 555 560
 10 Thr Trp Trp Arg Val His Asp Ala Met Pro Lys Lys Leu Gln Arg Phe
 565 570 575
 Cys Leu Leu Ser Ser Lys Gln Lys Ala Gln Leu Glu Trp Asp Arg Arg
 580 585 590
 15 Gln Ala Glu Lys Ala Asn Tyr Arg Asp Gly His Trp Arg Ile Lys Ile
 595 600 605
 Lys Asp Lys Arg Leu Thr Thr Cys Phe Glu Asp Phe Cys Phe Trp Glu
 610 615 620
 Ser Met Leu Trp His Trp Gly Glu Thr Gln Thr Asn Ser Thr Val Ala
 625 630 635 640
 20 Ala Asp Ser Ser Ser Thr Ala
 645

25 <210> 11
 <211> 1857
 <212> DNA
 <213> Arabidopsis thaliana

30 <220>
 <221> CDS
 <222> (1)..(1857)

<400> 11
 35 atg ata cct tcc tca agt ccc atg gag tca aga cat cga ctc tcg ttc 48
 Met Ile Pro Ser Ser Ser Pro Met Glu Ser Arg His Arg Leu Ser Phe
 1 5 10 15

40 tca aaa gag aag aca agt agg agg aga ttt caa aga att gag aag ggt 96
 Ser Lys Glu Lys Thr Ser Arg Arg Arg Phe Gln Arg Ile Glu Lys Gly
 20 25 30

45 gtc aag ttc aac act ctg aaa ctt gtg ttg att tgt ata atg ctt gga 144
 Val Lys Phe Asn Thr Leu Lys Leu Val Leu Ile Cys Ile Met Leu Gly
 35 40 45

50 gct ttg ttc acg atc tac cgt ttt cgt tat cca ccg cta caa att cct 192
 Ala Leu Phe Thr Ile Tyr Arg Phe Arg Tyr Pro Pro Leu Gln Ile Pro
 50 55 60

55 gaa att cca act agt ttt ggt ctt act act gat cct cgc tat gta gct 240
 Glu Ile Pro Thr Ser Phe Gly Leu Thr Thr Asp Pro Arg Tyr Val Ala
 65 70 75 80

55 aca gct gag atc aac tgg aac cat atg tca aat ctt gtt gag aag cac 288
 Thr Ala Glu Ile Asn Trp Asn His Met Ser Asn Leu Val Glu Lys His
 85 90 95

60 gta ttt ggt aga agc gag tat caa gga att ggt ctt ata aat ctt aac 336
 Val Phe Gly Arg Ser Glu Tyr Gln Gly Ile Gly Leu Ile Asn Leu Asn
 100 105 110

22

	gat aac gag att gat cga ttc aag gag gta acg aaa tct gac tgt gat	384
	Asp Asn Glu Ile Asp Arg Phe Lys Glu Val Thr Lys Ser Asp Cys Asp	
	115 120 125	
5	cat gta gct ttg cat cta gat tat gct gca aag aac ata aca tgg gaa	432
	His Val Ala Leu His Leu Asp Tyr Ala Ala Lys Asn Ile Thr Trp Glu	
	130 135 140	
10	tct tta tac ccg gaa tgg att gat gaa gtt gaa gaa ttc gaa gtc cct	480
	Ser Leu Tyr Pro Glu Trp Ile Asp Glu Val Glu Glu Phe Glu Val Pro	
	145 150 155 160	
15	act tgt cct tct ctg cct ttg att caa att cct ggc aag cct cgg att	528
	Thr Cys Pro Ser Leu Pro Leu Ile Gln Ile Pro Gly Lys Pro Arg Ile	
	165 170 175	
20	gat ctt gta att gcc aag ctt ccg tgt gat aaa tca gga aaa tgg tct	576
	Asp Leu Val Ile Ala Lys Leu Pro Cys Asp Lys Ser Gly Lys Trp Ser	
	180 185 190	
25	aga gat gtg gct cgc ttg cat tta caa ctt gca gca gct cga gtg gcg	624
	Arg Asp Val Ala Arg Leu His Leu Gln Leu Ala Ala Arg Val Ala	
	195 200 205	
30	gct tct tct aaa gga ctt cat aat gtt cat gtg att ttg gta tct gat	672
	Ala Ser Ser Lys Gly Leu His Asn Val His Val Ile Leu Val Ser Asp	
	210 215 220	
35	tgc ttt cca ata ccg aat ctt ttt acg ggt caa gaa ctt gtt gcc cgt	720
	Cys Phe Pro Ile Pro Asn Leu Phe Thr Gly Gln Glu Leu Val Ala Arg	
	225 230 235 240	
40	caa gga aac ata tgg ctg tat aag cct aat ctt cnc cag cta aga caa	768
	Gln Gly Asn Ile Trp Leu Tyr Lys Pro Asn Leu His Gln Leu Arg Gln	
	245 250 255	
45	aag tta cag ctt cct gtt ggt toc tgt gaa ctt tct gtt cct ctt caa	816
	Lys Leu Gln Leu Pro Val Gly Ser Cys Glu Leu Ser Val Pro Leu Gln	
	260 265 270	
50	gct aaa gat aat ttc tac tcc gca ggt gca aag aaa gaa gct tac gcg	864
	Ala Lys Asp Asn Phe Tyr Ser Ala Gly Ala Lys Lys Glu Ala Tyr Ala	
	275 280 285	
55	act atc ttg cat tct gcc caa ttt tat gtc tgt gga gcc att gca gct	912
	Thr Ile Leu His Ser Ala Gln Phe Tyr Val Cys Gly Ala Ile Ala Ala	
	290 295 300	
60	gca cag agc att cga atg tca ggc tct act cgt gat ctg gtc ata ctt	960
	Ala Gln Ser Ile Arg Met Ser Gly Ser Thr Arg Asp Leu Val Ile Leu	
	305 310 315 320	
65	gtt gat gaa acg ata agc gaa tac cat aaa agt ggc ttg gta gct gct	1008
	Val Asp Glu Thr Ile Ser Glu Tyr His Lys Ser Gly Leu Val Ala Ala	
	325 330 335	
70	gga tgg aag att caa atg ttt caa aga atc agg aac ccg aat gct gta	1056
	Gly Trp Lys Ile Gln Met Phe Gln Arg Ile Arg Asn Pro Asn Ala Val	
	340 345 350	
75	cca aat gcc tac aac gaa tgg aac tac agc aag ttt cgt ctt tgg caa	1104

23

	Pro Asn Ala Tyr Asn Glu Trp Asn Tyr Ser Lys Phe Arg Leu Trp Gln	
	355 360 365	
5	ctg act gaa tac agt aag atc atc ttc atc gat gca gac atg ctt atc Leu Thr Glu Tyr Ser Lys Ile Ile Phe Ile Asp Ala Asp Met Leu Ile	1152
	370 375 380	
10	ctg aga aac att gat ttc ctc ttc gag ttc cct gag ata tca gca act Leu Arg Asn Ile Asp Phe Leu Phe Glu Phe Pro Glu Ile Ser Ala Thr	1200
	385 390 395 400	
15	gga aac aat gct acg ctc ttc aac tct ggt cta atg gtg gtt gag cca Gly Asn Asn Ala Thr Leu Phe Asn Ser Gly Leu Met Val Val Glu Pro	1248
	405 410 415	
	tct aat tca aca ttc cag tta cta atg gat aac att aat gaa gtt gtg Ser Asn Ser Thr Phe Gln Leu Leu Met Asp Asn Ile Asn Glu Val Val	1296
	420 425 430	
20	tct tac aac gga gga gac caa ggt tac ctt aac gag ata ttc aca tgg Ser Tyr Asn Gly Gly Asp Gln Gly Tyr Leu Asn Glu Ile Phe Thr Trp	1344
	435 440 445	
25	tgg cat cgg att cca aaa cac atg aat ttc ttg aag cat ttc tgg gaa Trp His Arg Ile Pro Lys His Met Asn Phe Leu Lys His Phe Trp Glu	1392
	450 455 460	
30	gga gac gaa cct gag att aaa aaa atg aag acg agt cta ttt gga gct Gly Asp Glu Pro Glu Ile Lys Lys Met Lys Thr Ser Leu Phe Gly Ala	1440
	465 470 475 480	
35	gat cct cag atc cta tac gtt ctt cat tac cta ggt tat aac aaa ccc Asp Pro Pro Ile Leu Tyr Val Leu His Tyr Leu Gly Tyr Asn Lys Pro	1488
	485 490 495	
	tgg tta tgc ttc aga gac tat gac tgc aat tgg aat gtc gat att ttc Trp Leu Cys Phe Arg Asp Tyr Asp Cys Asn Trp Asn Val Asp Ile Phe	1536
	500 505 510	
40	cag gaa ttt gct agt gac gag gct cat aaa acc tgg tgg aga gtg cac Gln Glu Phe Ala Ser Asp Glu Ala His Lys Thr Trp Trp Arg Val His	1584
	515 520 525	
45	gac gca atg cct gaa aac ttg cat aag ttc tgt cta cta aga tgc aaa Asp Ala Met Pro Glu Asn Leu His Lys Phe Cys Leu Leu Arg Ser Lys	1632
	530 535 540	
50	cag aag gcg caa ctt gaa tgg gat agg aga caa gca gag aaa ggg aac Gln Lys Ala Gln Leu Glu Trp Asp Arg Arg Gln Ala Glu Lys Gly Asn	1680
	545 550 555 560	
55	tac aaa gat gga cat tgg aag ata aag atc aaa gac aag aga ctt aag Tyr Lys Asp Gly His Trp Lys Ile Lys Ile Lys Asp Lys Arg Leu Lys	1728
	565 570 575	
	act tgt ttc gaa gat ttc tgc ttt tgg gag agt atg ctt tgg cat tgg Thr Cys Phe Glu Asp Phe Cys Phe Trp Glu Ser Met Leu Trp His Trp	1776
	580 585 590	
60	ggt gag acg aac tct acc aac aat tot tcc acc acc acc act tca tca	1824
	Gly Glu Thr Asn Ser Thr Asn Asn Ser Ser Thr Thr Thr Thr Ser Ser	

24

595

600

605

ccg ccg cat aaa acc gct ctc cct tcc ctg tga
 Pro Pro His Lys Thr Ala Leu Pro Ser Leu
 610 615

1857

5

10

<210> 12
 <211> 618
 <212> PRT
 <213> Arabidopsis thaliana

15

20

25

30

35

40

45

50

55

60

<400> 12
 Met Ile Pro Ser Ser Ser Pro Met Glu Ser Arg His Arg Leu Ser Phe
 1 5 10 15
 Ser Lys Glu Lys Thr Ser Arg Arg Arg Phe Gln Arg Ile Glu Lys Gly
 20 25 30
 Val Lys Phe Asn Thr Leu Lys Leu Val Leu Ile Cys Ile Met Leu Gly
 35 40 45
 Ala Leu Phe Thr Ile Tyr Arg Phe Arg Tyr Pro Pro Leu Gln Ile Pro
 50 55 60
 Glu Ile Pro Thr Ser Phe Gly Leu Thr Thr Asp Pro Arg Tyr Val Ala
 65 70 75 80
 Thr Ala Glu Ile Asn Trp Asn His Met Ser Asn Leu Val Glu Lys His
 85 90 95
 Val Phe Gly Arg Ser Glu Tyr Gln Gly Ile Gly Leu Ile Asn Leu Asn
 100 105 110
 Asp Asn Glu Ile Asp Arg Phe Lys Glu Val Thr Lys Ser Asp Cys Asp
 115 120 125
 His Val Ala Leu His Leu Asp Tyr Ala Ala Lys Asn Ile Thr Trp Glu
 130 135 140
 Ser Leu Tyr Pro Glu Trp Ile Asp Glu Val Glu Glu Phe Glu Val Pro
 145 150 155 160
 Thr Cys Pro Ser Leu Pro Leu Ile Gln Ile Pro Gly Lys Pro Arg Ile
 165 170 175
 Asp Leu Val Ile Ala Lys Leu Pro Cys Asp Lys Ser Gly Lys Trp Ser
 180 185 190
 Arg Asp Val Ala Arg Leu His Leu Gln Leu Ala Ala Arg Val Ala
 195 200 205
 Ala Ser Ser Lys Gly Leu His Asn Val His Val Ile Leu Val Ser Asp
 210 215 220
 Cys Phe Pro Ile Pro Asn Leu Phe Thr Gly Gln Glu Leu Val Ala Arg
 225 230 235 240
 Gln Gly Asn Ile Trp Leu Tyr Lys Pro Asn Leu His Gln Leu Arg Gln
 245 250 255
 Lys Leu Gln Leu Pro Val Gly Ser Cys Glu Leu Ser Val Pro Leu Gln
 260 265 270
 Ala Lys Asp Asn Phe Tyr Ser Ala Gly Ala Lys Lys Glu Ala Tyr Ala
 275 280 285
 Thr Ile Leu His Ser Ala Gln Phe Tyr Val Cys Gly Ala Ile Ala Ala
 290 295 300
 Ala Gln Ser Ile Arg Met Ser Gly Ser Thr Arg Asp Leu Val Ile Leu
 305 310 315 320
 Val Asp Glu Thr Ile Ser Glu Tyr His Lys Ser Gly Leu Val Ala Ala
 325 330 335
 Gly Trp Lys Ile Gln Met Phe Gln Arg Ile Arg Asn Pro Asn Ala Val
 340 345 350
 Pro Asn Ala Tyr Asn Glu Trp Asn Tyr Ser Lys Phe Arg Leu Trp Gln
 355 360 365
 Leu Thr Glu Tyr Ser Lys Ile Ile Phe Ile Asp Ala Asp Met Leu Ile
 370 375 380

25

Leu Arg Asn Ile Asp Phe Leu Phe Glu Phe Pro Glu Ile Ser Ala Thr
 385 390 395 400
 Gly Asn Asn Ala Thr Leu Phe Asn Ser Gly Leu Met Val Val Glu Pro
 405 410 415
 5 Ser Asn Ser Thr Phe Gln Leu Leu Met Asp Asn Ile Asn Glu Val Val
 420 425 430
 Ser Tyr Asn Gly Gly Asp Gln Gly Tyr Leu Asn Glu Ile Phe Thr Trp
 435 440 445
 10 Trp His Arg Ile Pro Lys His Met Asn Phe Leu Lys His Phe Trp Glu
 450 455 460
 Gly Asp Glu Pro Glu Ile Lys Lys Met Lys Thr Ser Leu Phe Gly Ala
 465 470 475 480
 Asp Pro Pro Ile Leu Tyr Val Leu His Tyr Leu Gly Tyr Asn Lys Pro
 485 490 495
 15 Trp Leu Cys Phe Arg Asp Tyr Asp Cys Asn Trp Asn Val Asp Ile Phe
 500 505 510
 Gln Glu Phe Ala Ser Asp Glu Ala His Lys Thr Trp Trp Arg Val His
 515 520 525
 20 Asp Ala Met Pro Glu Asn Leu His Lys Phe Cys Leu Leu Arg Ser Lys
 530 535 540
 Gln Lys Ala Gln Leu Glu Trp Asp Arg Arg Gln Ala Glu Lys Gly Asn
 545 550 555 560
 Tyr Lys Asp Gly His Trp Lys Ile Lys Ile Lys Asp Lys Arg Leu Lys
 565 570 575
 25 Thr Cys Phe Glu Asp Phe Cys Phe Trp Glu Ser Met Leu Trp His Trp
 580 585 590
 Gly Glu Thr Asn Ser Thr Asn Asn Ser Ser Thr Thr Thr Ser Ser
 595 600 605
 30 Pro Pro His Lys Thr Ala Leu Pro Ser Leu
 610 615

35 <210> 13
 <211> 2130
 <212> DNA
 <213> Arabidopsis thaliana

40 <220>
 <221> CDS
 <222> (1)..(1980)

45 <400> 13
 atg gca aac tct ccc gct gct cct gca ccc acc acc aca acc ggt ggt 48
 Met Ala Asn Ser Pro Ala Ala Pro Ala Pro Thr Thr Thr Thr Gly Gly
 1 5 10 15
 50 gac tcc cgg cga cgc ctg tcc gcg tcc ata gaa gca ata tgc aag agg 96
 Asp Ser Arg Arg Arg Leu Ser Ala Ser Ile Glu Ala Ile Cys Lys Arg
 20 25 30
 aga ttc cgg aga aat agc aaa gga ggt ggc aga tcg gat atg gtg aaa 144
 Arg Phe Arg Arg Asn Ser Lys Gly Gly Arg Ser Asp Met Val Lys
 35 40 45
 55 ccg ttt aat atc ata aat ttt tcg aca caa gac aaa aac agt agt tgt 192
 Pro Phe Asn Ile Ile Asn Phe Ser Thr Gln Asp Lys Asn Ser Ser Cys
 50 55 60
 60 tgt tgt ttc acc aag ttt cag atc gtg aag ctt ctc ttg ttt atc ctt 240
 Cys Cys Phe Thr Lys Phe Gln Ile Val Lys Leu Leu Leu Phe Ile Leu
 65 70 75 80

26

5	ctc tct gcc act ctc ttc acc att atc tat tct cct gaa gct tat cat	288
	Leu Ser Ala Thr 85 Leu Phe Thr Ile Ile Tyr Ser Pro Glu Ala Tyr His 95	
10	cat tct ctt tcc cac tca tct tct cga tgg ata tgg aga aga caa gat	336
	His Ser Leu Ser 100 His Ser Ser Ser Arg Trp Ile Trp Arg Arg Glu Asp 110	
15	cca cgt tac ttc tcg gat ctg gat ata aac tgg gac gat gtg act aaa	384
	Pro Arg Tyr Phe Ser Asp Leu Asp Ile Asn Trp Asp Asp Val Thr Lys 115 120 125	
20	acc ctt gag aac atc gaa gaa ggc cgt acg atc ggt gtc ttg aat ttt	432
	Thr Leu Glu Asn Ile Glu Glu Gly Arg Thr Ile Gly Val Leu Asn Phe 130 135 140	
25	gat tcg aac gag atc caa cga tgg aga gaa gta tcc aag agc aag gac	480
	Asp Ser Asn Glu Ile Gln Arg Trp Arg Glu Val Ser Lys Ser Lys Asp 145 150 155 160	
30	aat ggg gat gaa gaa aaa gtt gtt gta ttg aat cta gat tac gca gac	528
	Asn Gly Asp Glu Glu Lys Val Val Val Leu Asn Leu Asp Tyr Ala Asp 165 170 175	
35	aag aat gtg act tgg gac gca cta tat cca gag tgg atc gat gag gag	576
	Lys Asn Val Thr Trp Asp Ala Leu Tyr Pro Glu Trp Ile Asp Glu Glu 180 185 190	
40	caa gaa aca gag gtc cct gtt tgt cct aat atc ccg aac att aag gta	624
	Gln Glu Thr Glu Val Pro Val Cys Pro Asn Ile Pro Asn Ile Lys Val 195 200 205	
45	cct aca aga aga ctc gat ctg atc gtc gtg aaa ctt cct tgt cgg aaa	672
	Pro Thr Arg Arg Leu Asp Leu Ile Val Val Lys Leu Pro Cys Arg Lys 210 215 220	
50	gaa ggg aat tgg tcg aga gac gtc ggg aga ttg cat cta cag cta gcg	720
	Glu Gly Asn Trp Ser Arg Asp Val Gly Arg Leu His Leu Gln Leu Ala 225 230 235 240	
55	gct gca act gtg gcg gct tcg gcc aaa ggg ttt ttc agg gga cat gtg	768
	Ala Ala Thr Val Ala Ala Ser Ala Lys Gly Phe Phe Arg Gly His Val 245 250 255	
60	ttt ttt gta tct aga tgc ttt ccg att ccg aat ctt ttc cgg tgt aaa	816
	Phe Phe Val Ser Arg Cys Phe Pro Ile Pro Asn Leu Phe Arg Cys Lys 260 265 270	
65	gat ctt gtg tct cgg aga ggc gat gtt tgg ttg tac aaa cct aat ctt	864
	Asp Leu Val Ser Arg Arg Gly Asp Val Trp Leu Tyr Lys Pro Asn Leu 275 280 285	
70	gat acc ttg aga gac aag ctt cag ctg cct gta ggg tct tgt gag cta	912
	Asp Thr Leu Arg Asp Lys Leu Gln Leu Pro Val Gly Ser Cys Glu Leu 290 295 300	
75	tct ctt cct ctt ggc atc caa gat agg cca ago tta gga aac cct aaa	960
	Ser Leu Pro Leu Gly Ile Gln Asp Arg Pro Ser Leu Gly Asn Pro Lys 305 310 315 320	
aga gaa gct tac gca aca att ctt cac tca gct cac gtt tac gtc tgc		1008

27

Arg Glu Ala Tyr Ala Thr Ile Leu His Ser Ala His Val Tyr Val Cys

325

330

335

5 ggt gca atc gcc gcg gct cag agc ata aga cag tct ggt tcg acg aga 1056
Gly Ala Ile Ala Ala Ala Gln Ser Ile Arg Gln Ser Gly Ser Thr Arg
340 345 350

10 gac ctt gtt atc ctt gtt gat gac aac atc agc ggt tac cac cgg agt 1104
Asp Leu Val Ile Leu Val Asp Asp Asn Ile Ser Gly Tyr His Arg Ser
355 360 365

15 gga cta gaa gcc gcg ggt tgg caa atc cgg acg ata cag agg att cga 1152
Gly Leu Glu Ala Ala Gly Trp Gln Ile Arg Thr Ile Gln Arg Ile Arg
370 375 380

20 aac cct aag gca gag aaa gat gct tac aac gaa tgg aac tac agc aag 1200
Asn Pro Lys Ala Glu Lys Asp Ala Tyr Asn Glu Trp Asn Tyr Ser Lys
385 390 395 400

ttc cgg cta tgg cag ctg act gat tac gac aaa atc att ttc atc gac 1248
Phe Arg Leu Trp Gln Leu Thr Asp Tyr Asp Lys Ile Ile Phe Ile Asp
405 410 415

25 gcg gat ctc tta atc ttg aga aac atc gat ttc ttg ttc tcg atg cct 1296
Ala Asp Leu Leu Ile Leu Arg Asn Ile Asp Phe Leu Phe Ser Met Pro
420 425 430

30 gag atc tca gct aca gga aac aat gga act ctg ttt aat tca gga gtt 1344
Glu Ile Ser Ala Thr Gly Asn Asn Gly Thr Leu Phe Asn Ser Gly Val
435 440 445

35 atg gtg atc gag cct tgc aac tgt acg ttt cag ctt ctg atg gaa cat 1392
Met Val Ile Glu Pro Cys Asn Cys Thr Phe Gln Leu Leu Met Glu His
450 455 460

40 ata aac gag att gag tct tat aac ggt gga gat caa ggt tac tta aac 1440
Ile Asn Glu Ile Glu Ser Tyr Asn Gly Gly Asp Gln Gly Tyr Leu Asn
465 470 475 480

gag gta ttc aca tgg tgg cac cgg att cca aaa cat atg aat ttc ttg 1488
Glu Val Phe Thr Trp Trp His Arg Ile Pro Lys His Met Asn Phe Leu
485 490 495

45 aag cat ttt tgg att ggc gat gaa gat gac gcg aaa cgc aag aaa aca 1536
Lys His Phe Trp Ile Gly Asp Glu Asp Asp Ala Lys Arg Lys Lys Thr
500 505 510

50 gag ctt ttt gga gca gag cct cct gtt ctt tat gtt ctt cat tac ctt 1584
Glu Leu Phe Gly Ala Glu Pro Pro Val Leu Tyr Val Leu His Tyr Leu
515 520 525

55 ggg atg aag ccg tgg tta tgt tac cgt gac tac gac tgt aac ttc aac 1632
Gly Met Lys Pro Trp Leu Cys Tyr Arg Asp Tyr Asp Cys Asn Phe Asn
530 535 540

60 tcc gac ata ttc gtt gag ttt gct acc gat atc gct cat cga aaa tgg 1680
Ser Asp Ile Phe Val Glu Phe Ala Thr Asp Ile Ala His Arg Lys Trp
545 550 555 560

trg atg gtc cac gac gcc atg cca cag gaa ctt cac caa ttc tgt tac 1728
Trp Met Val His Asp Ala Met Pro Gln Glu Leu His Gln Phe Cys Tyr

575

	ttg cga tcc aag caa aag gca cag ctg gaa tat gat cgc cgg caa gca	1776
5	Leu Arg Ser Lys Gln Lys Ala Gln Leu Glu Tyr Asp Arg Arg Gln Ala	
	580 585 590	
	gag gcc gca aat tat gcc gac ggt cat tgg aaa ata aga gta aag gac	1824
	Glu Ala Ala Asn Tyr Ala Asp Gly His Trp Lys Ile Arg Val Lys Asp	
	595 600 605	
10	ccg aga ttc aaa att tgc atc gac aaa tta tgt aat tgg aaa agt atg	1872
	Pro Arg Phe Lys Ile Cys Ile Asp Lys Leu Cys Asn Trp Lys Ser Met	
	610 615 620	
15	ctg cgg cat tgg ggc gaa tca aat tgg act gac tac gag tct ttt gtt	1920
	Leu Arg His Trp Gly Glu Ser Asn Trp Thr Asp Tyr Glu Ser Phe Val	
	625 630 635 640	
20	ccc acc cca cca gcc att acc gta gac cgg aga tca tca ctt ccc ggc	1968
	Pro Thr Pro Pro Ala Ile Thr Val Asp Arg Arg Ser Ser Leu Pro Gly	
	645 650 655	
	cat aac ttg tga cgcaataatt atacatactt attaatggat ttcattgagtt	2020
25	His Asn Leu	
	660	
	tttttggtttg aattgtttgct gegagattag gtgaatatca gttgtgtaac tatatctttt	2080
	tcctatagtt tgttcaaatt gaataaaaaca tttttttgca gtttaaccac	2130
30		
	<210> 14	
	<211> 659	
	<212> PRT	
35	<213> Arabidopsis thaliana	
	<400> 14	
	Met Ala Asn Ser Pro Ala Ala Pro Ala Pro Thr Thr Thr Thr Gly Gly	
	1 5 10 15	
40	Asp Ser Arg Arg Arg Leu Ser Ala Ser Ile Glu Ala Ile Cys Lys Arg	
	20 25 30	
	Arg Phe Arg Arg Asn Ser Lys Gly Gly Arg Ser Asp Met Val Lys	
	35 40 45	
45	Pro Phe Asn Ile Ile Asn Phe Ser Thr Gln Asp Lys Asn Ser Ser Cys	
	50 55 60	
	Cys Cys Phe Thr Lys Phe Gln Ile Val Lys Leu Leu Phe Ile Leu	
	65 70 75 80	
	Leu Ser Ala Thr Leu Phe Thr Ile Ile Tyr Ser Pro Glu Ala Tyr His	
	85 90 95	
50	His Ser Leu Ser His Ser Ser Ser Arg Trp Ile Trp Arg Arg Gln Asp	
	100 105 110	
	Pro Arg Tyr Phe Ser Asp Leu Asp Ile Asn Trp Asp Asp Val Thr Lys	
	115 120 125	
55	Thr Leu Glu Asn Ile Glu Glu Gly Arg Thr Ile Gly Val Leu Asn Phe	
	130 135 140	
	Asp Ser Asn Glu Ile Gln Arg Trp Arg Glu Val Ser Lys Ser Lys Asp	
	145 150 155 160	
	Asn Gly Asp Glu Glu Lys Val Val Val Leu Asn Leu Asp Tyr Ala Asp	
	165 170 175	
60	Lys Asn Val Thr Trp Asp Ala Leu Tyr Pro Glu Trp Ile Asp Glu Glu	
	180 185 190	
	Gln Glu Thr Glu Val Pro Val Cys Pro Asn Ile Pro Asn Ile Lys Val	

29

195 200 205
 Pro Thr Arg Arg Leu Asp Leu Ile Val Val Lys Leu Pro Cys Arg Lys
 210 215 220
 5 Glu Gly Asn Trp Ser Arg Asp Val Gly Arg Leu His Leu Gln Leu Ala
 225 230 235 240
 Ala Ala Thr Val Ala Ala Ser Ala Lys Gly Phe Phe Arg Gly His Val
 245 250 255
 Phe Phe Val Ser Arg Cys Phe Pro Ile Pro Asn Leu Phe Arg Cys Lys
 260 265 270
 10 Asp Leu Val Ser Arg Arg Gly Asp Val Trp Leu Tyr Lys Pro Asn Leu
 275 280 285
 Asp Thr Leu Arg Asp Lys Leu Gln Leu Pro Val Gly Ser Cys Glu Leu
 290 295 300
 15 Ser Leu Pro Leu Gly Ile Gln Asp Arg Pro Ser Leu Gly Asn Pro Lys
 305 310 315 320
 Arg Glu Ala Tyr Ala Thr Ile Leu His Ser Ala His Val Tyr Val Cys
 325 330 335
 Gly Ala Ile Ala Ala Ala Gln Ser Ile Arg Gln Ser Gly Ser Thr Arg
 340 345 350
 20 Asp Leu Val Ile Leu Val Asp Asp Asn Ile Ser Gly Tyr His Arg Ser
 355 360 365
 Gly Leu Glu Ala Ala Gly Trp Gln Ile Arg Thr Ile Gln Arg Ile Arg
 370 375 380
 25 Asn Pro Lys Ala Glu Lys Asp Ala Tyr Asn Glu Trp Asn Tyr Ser Lys
 385 390 395 400
 Phe Arg Leu Trp Gln Leu Thr Asp Tyr Asp Lys Ile Ile Phe Ile Asp
 405 410 415
 Ala Asp Leu Leu Ile Leu Arg Asn Ile Asp Phe Leu Phe Ser Met Pro
 420 425 430
 30 Glu Ile Ser Ala Thr Gly Asn Asn Gly Thr Leu Phe Asn Ser Gly Val
 435 440 445
 Met Val Ile Glu Pro Cys Asn Cys Thr Phe Gln Leu Leu Met Glu His
 450 455 460
 35 Ile Asn Glu Ile Glu Ser Tyr Asn Gly Gly Asp Gln Gly Tyr Leu Asn
 465 470 475 480
 Glu Val Phe Thr Trp Trp His Arg Ile Pro Lys His Met Asn Phe Leu
 485 490 495
 Lys His Phe Trp Ile Gly Asp Glu Asp Asp Ala Lys Arg Lys Lys Thr
 500 505 510
 40 Glu Leu Phe Gly Ala Glu Pro Pro Val Leu Tyr Val Leu His Tyr Leu
 515 520 525
 Gly Met Lys Pro Trp Leu Cys Tyr Arg Asp Tyr Asp Cys Asn Phe Asn
 530 535 540
 45 Ser Asp Ile Phe Val Glu Phe Ala Thr Asp Ile Ala His Arg Lys Trp
 545 550 555 560
 Trp Met Val His Asp Ala Met Pro Gln Glu Leu His Gln Phe Cys Tyr
 565 570 575
 Leu Arg Ser Lys Gln Lys Ala Gln Leu Glu Tyr Asp Arg Arg Gln Ala
 580 585 590
 50 Glu Ala Ala Asn Tyr Ala Asp Gly His Trp Lys Ile Arg Val Lys Asp
 595 600 605
 Pro Arg Phe Lys Ile Cys Ile Asp Lys Leu Cys Asn Trp Lys Ser Met
 610 615 620
 55 Leu Arg His Trp Gly Glu Ser Asn Trp Thr Asp Tyr Glu Ser Phe Val
 625 630 635 640
 Pro Thr Pro Pro Ala Ile Thr Val Asp Arg Arg Ser Ser Leu Pro Gly
 645 650 655
 His Asn Leu

60

<210> 15

<211> 15294
<212> DNA
<213> Artificial Sequence

5 <220>

<220>
<223> Description of Artificial Sequence: vector

10 <400> 15

ggcggggagg	gttcgagaag	ggggggcacc	ccccttcggc	gtgcgagggtc	acgcgcacag	60	
ggcgcagccc	tgggttaaaaa	caaggtttat	aaatatgggt	ttaaaagcag	gttaaaagac	120	
aggttagcgg	tggccgaaaa	acgggaggaa	acccttgcaa	atgctggatt	ttctgcctgt	180	
ggacagcccc	tcaaattgtca	ataggtgcgc	ccctcatctg	tcagcactct	gcccccaag	240	
15	tgtcaaggat	cgcgcgcctc	atctgtcagt	agtcgcgcgc	ctcaagtgtc	aataccgcag	300
ggcacttatt	cccaggcttg	tccacatcat	ctgtgggaaa	ctcgcgtaaa	atcaggcggt	360	
ttcgccgatt	gcccagcttg	gccagctcca	cgtcgcgcgc	cgaatcgag	cctgccccctc	420	
atctgtcaac	gcccgcgcgc	gtgagtcggc	ccctcaagt	tcaacgtccg	ccctcatct	480	
gtcagtgagg	gccaagtgtt	ccgcgaggta	tccacaacgc	ggcgcgcgc	ggtgtctcgc	540	
20	acacggcttc	gacggcggtt	ctggcgcggt	tgcaggggcca	tagacgggcg	ccagccacgc	600
ggcgagggca	accagcccgc	tgagcgtcgc	aaaggcgctc	ggtcttgcc	tgctcgtcgc	660	
tgatgacatt	caccagctcc	gcgaagtcgc	tcttcttgat	ggagcgcag	gggacgtgct	720	
tggcaatcac	gcgcaccccc	cggcggtttt	agcggctaaa	aaagtoatgg	ctctgcccc	780	
ggcgaggacca	cgcctcatcat	gaccttgcca	agctcgtcct	gcttctcttc	gatcttcgcc	840	
25	agcagggcga	ggatcgtggc	atcacccaac	cgcgcgcgtc	gggggtcgtc	ggtgagccag	900
agtttcagca	ggcgcgcag	gccccccagg	tcgccattga	tgcggggccag	ctcgcggacg	960	
tgctcatagt	ccacgacgcc	cgtgattttg	tagccctggc	cgcgcggccag	caggtaggcc	1020	
gacaggctca	tgcgcggcgc	cgcgcgcctt	tcctcaatcg	ctcttcgttc	gtctggaagg	1080	
cagtacacct	tgataggtgg	gctgccttcc	ctggttggct	tggtttcatc	agccatccgc	1140	
30	ttgccctcat	ctgttacgcc	ggcggtagcc	ggccagcctc	gcagagcagg	attcccgttg	1200
agcaccgccca	ggtgcgaata	agggacagtg	aagaaggaa	acccgctcgc	gggtgggcct	1260	
acttcacctc	tcctgcccgc	ctgacgcctg	tggatacacc	aaggaaagtc	tacacgaacc	1320	
ctttggcaaa	atcctgtata	tcgtgcgaaa	aaggatggat	ataccgaaaa	aatcgctata	1380	
atgaccccca	agcaggggtta	tgcagcggaa	aagcggcag	otccccgaag	ggagaaaggc	1440	
35	ggcagcgtat	ccggtaaagc	gcagggtag	aacaggagag	cgcacgaggg	agcttccagg	1500
gggaaacgcc	tgggtatctt	atagtcctgt	cgggtttcgc	cacctctgac	ttgagcgtcg	1560	
atctttgtga	tgctcgtcag	gggggaggag	cctatggaaa	aacgcagca	acgcggcctt	1620	
tttacgggtc	ctggcctttt	gctggccttt	tgctcacatg	ttctttcctg	cgttatcccc	1680	
tgattctgtg	gataaccgta	ttaccgcctt	tgagtgaagt	gataccgctc	gcccagccgc	1740	
40	aacgacggag	cgcagcgagt	cagtgaagca	ggaagcggaa	gagcgcacga	aggccgcctc	1800
agagggccag	cgcgcgcgtg	aggctcggac	gctagggcag	ggcatgaaaa	agcccgtagc	1860	
gggctgctac	ggggtctgta	cgcgggtgaa	agggggaggg	gatgttgtct	acatggctct	1920	
gctgtagtga	gtgggttgcg	ctccggcagc	ggtcctgac	aatcgtcacc	ctttctcggc	1980	
ccttcaacgt	tcctgacaa	gagcctcctt	ttcgccaatc	catcgacaat	caccgcgagt	2040	
45	ccctgctcga	acgctgcgtc	cggacccggt	tcgtcgaagg	cgtctatcgc	ggcccgcaac	2100
agcggcgaga	gcggagcctg	tccaacgggt	ccgcgcgcgt	cgcgcgcctc	gctgtcgcgc	2160	
gcctgctcct	caagcacggc	cccaacagtg	aagttagctga	ttgtcatcag	cgcattgacg	2220	
gcgtccccgc	ccgaaaaacc	cgcctcgcag	aggaagcgaa	gctgcgcgtc	ggccggtttcc	2280	
atctgcgggtg	cgcgcggctg	cgtgcgcgca	tggatgcgcg	cgcctatcgc	gtaggcgagc	2340	
50	agcgcctgce	tgaagctgcg	ggcattcccg	atcagaaatg	agcgcagctc	gtcgtcggct	2400
ctcggcacgc	aatgcgtatg	attctcgcgc	agcatggctt	cggccagtg	gtcagagcag	2460	
gcccgcctgt	tctgaagtg	ccagtaagc	gcccgtcgt	gaacccccaa	ccgttccgc	2520	
agtttgcgtg	tcgtcagacc	gtctacgcgc	acctcgttca	acaggtccag	ggcggcagcg	2580	
atcactgtat	tgggtgcgaa	ccttgcctat	cctgacactt	tatcaactgat	aaacataata	2640	
55	tgtccacca	cttatcagtg	ataaagaatc	cgcgcgttca	atcggaccag	cggaggctgg	2700
tcgggaggcc	agacgtgaaa	cccaacatac	ccctgatcgt	aattctgagc	actgtcgcgc	2760	
tcgacgctgt	cggcatcggc	ctgattatgc	cgggtcgtgc	gggcctcctg	cgcgatctgg	2820	
ttcactcgaa	cgcgtcacc	gcccactatg	gcattctgct	ggcgtctgat	gcgttgggtc	2880	
60	aatttgcctg	cgcacctgtg	ctgggcgcgc	tgtcggatcg	tttcgggcgc	cggccaatct	2940
tgctcgtctc	gctggccggc	gccaagatct	ggggaacct	gtggttggca	tgcacataca	3000	
aatggacgaa	cggataaacc	ttttcacgcc	cttttaata	tcgattatt	ctaataaacg	3060	
ctctttctct	ttaggtttac	ccgccaatat	atcctgtcaa	acactgatag	tttaactga	3120	

31

	aggcgggaaa	cgacaatctg	atcatgagcg	gagaattaag	ggagtcacgt	tatgaccccc	3180
	gccgatgacg	cgggacaagc	cgttttacgt	ttggaactga	cagaaccgca	acgttgaagg	3240
	agccactcag	cggatctgaa	ttcccgatct	agtaacatag	atgacaccgc	gcgcgataat	3300
5	ttatcctagt	ttgcgcgcta	tattttgttt	tctatcgctg	attaaatgta	taattgctgg	3360
	actctaataa	taaaaaacca	tctcataaat	aacgtcatgc	attacatggt	aattattaca	3420
	tgettaacgt	aattcaacag	aaattatatg	ataatcatcg	caagaccggc	aacaggattc	3480
	aatcttaaga	aacttttattg	ccaaatgttt	gaacgatcgg	ggaaattcga	gctcggatcc	3540
	atcatgttac	aaactttttt	gctgtgagca	gtagatatgg	aaaccoggag	gacctaaagt	3600
10	atctgataga	taaagcacat	agcttggggt	tacagggtct	gggtggatgta	gttcacagtc	3660
	atgcaagcaa	taatgccact	gatggcctca	atggctttga	tattggccaa	ggttctcaag	3720
	aatcctactt	tcatgctgga	gagcaagggt	accataagtt	gtgggatagc	aggctgttca	3780
	actatgccaa	ttgggaggtt	cttcgtttcc	ttctttccaa	cttgaggtgg	tggctagaag	3840
	agtataactt	tgacggattt	cgatttgatg	gaataacttc	tatgctgtat	gttcacatg	3900
15	gaatcaatat	gggattttaca	ggaaactata	atgagtattt	cagcgaggct	acagatgttg	3960
	atgctgtggt	ctatttaattg	ttggccaata	atctgattca	caagattttc	ccagacgcaa	4020
	ctgttattgc	cgaagatggt	tctggatatg	cgggccttgg	ccggcctggt	tctgagggag	4080
	gaattgggtt	tgattacgcg	ctggcaatgg	caatcccaga	taagtggata	gattatttaa	4140
	agaataaaaa	tgatgaagat	tggcccatga	aggaagtaac	atcgagtttg	acaaatagga	4200
20	gatatacaga	gaagtgtata	gcatatgcgg	agagccatga	tcagtctatt	gtcggtgaca	4260
	agaccattgc	atcttctcta	atggacaaaag	agatgtattc	tgcatgtctt	tgcttgacag	4320
	atgcttctcc	tgttattgat	cgaggaattg	cgttccacaa	gatgatccat	ttttccacaa	4380
	tggtccttgg	aggagagggg	tacetcgaat	tcatgggtaa	cgagtttggc	catcctgagt	4440
	ggattgactt	ccctagagag	ggcaataatt	ggtgttatga	caaagttaga	cgccagtggg	4500
25	accttgccga	tagcgaacac	ttgagataca	agtttatgaa	tgcatttgat	agagctatga	4560
	attcgcctga	tgaaaagtgc	tcattcctcg	catcaggaaa	acagatagta	agcagcatgg	4620
	atgatgagaa	gaaggttggt	gtgtttgaac	gtggtgacct	ggtattttgt	ttcaacttcc	4680
	acccaaataa	cacatacga	gggtataaag	ttggatgtga	cttcccaggg	aagtacagag	4740
	ttgcactgga	cagtgtgct	tgsgaatttg	gtggccatgg	aagagctggg	catgatgttg	4800
30	accatttcac	atcaccagaa	ggaatacctg	gagttccaga	aacaaatttc	aatggctcgt	4860
	caaattccct	caaagtgcgt	tctcctgcgc	gaacatgtgt	ggcttattac	agagttgacg	4920
	aacgcattgc	agaaactgaa	gtttaccaga	cagacatttc	tagtgagcta	ctaccaacag	4980
	ccaatatcga	ggagagtga	gagaaactta	aagattcgtt	atctacaaat	atcagtaacg	5040
	ttgacgaact	catgtcagaa	actgaagtgt	accagacaga	catctctagt	gagctactac	5100
35	caacagccag	tatcgaggag	agtgcagaga	accctaaaga	ttcattatct	acaaatatca	5160
	gtacgtgggt	atcattggat	gtgggattcc	cgcctcttta	attatggaaa	ctgggaggta	5220
	cttaggtatc	ttctctcaaa	tgcgagatgg	tggttgatg	agttcaaaat	tgatggattt	5280
	agattcgatg	gtgtgacatc	aataatgtat	actcaccacg	gattatcggt	gggattcact	5340
	gggaactaca	aggaatactt	tggaactcgca	actgatgtgg	atgctgttgt	gtatctgatg	5400
40	ctggtoaacg	atcttattca	tggtcttttc	cagatgcaat	taccattggg	gaagatgtta	5460
	gcggaatgcc	gacattttgt	attcccgttc	aagatggggg	tggtggcttt	gactatcggc	5520
	tgcatatggc	aattgctgat	aaatggattg	agttgctcaa	gaaacgggat	gaggattgga	5580
	gagtgggtga	tattgttcat	acactgacaa	atagaagatg	gtoggaaaag	tgtgtttcat	5640
	acgctgaaag	tcatgatcaa	gctctagtcg	gtgacaaaac	tatagcattc	tggtgatggg	5700
45	acaaggatat	gtatgatatt	atggtctcgg	atagaccatc	aacatcatta	atagatcgtg	5760
	ggatagcatt	gcacaagatg	attaggcttg	taactatggg	attaggagga	gaagggtacc	5820
	taaatttcat	gggaaatgaa	ttcgccacc	ctgagtggtg	tgatttccct	agggctgaac	5880
	aacacctctc	tgatggctca	gtaattcccg	gaaaccaatt	cagttatgat	aaatgcagac	5940
	ggagatttga	cotgggagat	gcagaatatt	taagataccg	tggttgcaaa	gaatttgacc	6000
50	gggctatgca	gtatcttgaa	gataaatatg	agtttatgac	ttcagaacac	cagttcatat	6060
	cacgaaagga	tgaaggagat	aggatgattg	tatttgaaaa	aggaaacctt	gtttttgtct	6120
	ttaattttca	ctggacaaaa	agctattcag	actatcgcat	aggctgcctg	aagcctggaa	6180
	aatacaaggt	tgcttctggc	tcagatgac	cacttcttgg	tggtctcggg	agaattgatc	6240
	ataatgccga	atatttccac	tttgaggatc	ggtatgagta	tcgtcctcgt	tcaattatgg	6300
55	tgtatgcacc	tagtagaaca	gcagtgggtc	atgcactagt	agacaaagaa	gaagaagaag	6360
	aagaagaagt	agcagtagta	gaagaagtag	tagtagaaga	agaatgaacg	aacttgtgat	6420
	cgcgttgaaa	gattttgaagg	ctacatagct	ctagagtcga	cctgcattga	atcagaaata	6480
	attggaggag	atgagtaaaa	gttaccactt	gttgagctgt	gtgagtgagt	gagtgagaat	6540
	gaggaggtgc	ctgccttatt	tgtagcaggt	ttcagtgaca	cgtgtcaaga	gaatagcggg	6600
60	tggctatccc	ttagcagaag	gcaactgtgg	acactgtatt	atagggaaat	gctcatcgac	6660
	agtattatgg	gcccctctct	tggtgattca	cggctgggact	tcaacttggg	ccttgcaagt	6720
	ggccctgccc	gttctgtctc	ctagtatcta	aaaaactaaa	ccaactccct	cctaccgcta	6780
	ccacttgaca	ttcctatgtc	tctgtttaat	taaatatta	ttatagtaat	taaaaaaat	6840

	atctagggtac	tgggtactggt	ccctccctcc	actagaatat	tagttacttc	ccccttagct	6900
	ttgtattcca	aattactgta	aataattttt	ctaatttttt	acgacaaaca	agatctaatt	6960
	atgaatgcac	aattcttaag	gttgaataca	ttactttact	tgggttagcc	tatatataag	7020
	tgcatttttag	tattaagatt	gagatgcatg	gttctattac	aaaattgata	cactgctaaa	7080
5	ggaaggatgg	ttaaaaacaa	cattcaatgt	ttgttacatt	tcttctctatt	gtattttttt	7140
	tttaacgagc	ttcccggtata	catcataaca	tgtctccggt	ccacttggca	ggaaaaaaa	7200
	atacccaaac	aggaagatag	tgtcaagtat	atccatagat	gaggacttaa	tggataggct	7260
	tttcgaggat	tcataaatca	taatatctgg	cggaggagtc	aattaaatac	ttgtggtttg	7320
	tatcctgatt	actccgtcaa	cagccaaata	gaaaagtgtg	aaaagagaga	aaggatttgg	7380
10	tacaagatac	tgttgcattt	gttaagtaat	gaacaaaacg	gagtaacata	attttctatc	7440
	tcgttaaagc	ttcacgctgc	cgaagcact	caggcgcaa	gggctgctaa	ggaagcggaa	7500
	cacgtagaaa	gccagtcgc	agaaacggtg	ctgaccccg	atgaatgtca	gctactgggc	7560
	tatctggaca	agggaaaacg	caagcgcaaa	gagaaagcag	gtagcttgca	gtgggcttac	7620
	atggcgatag	ctagactggg	cggttttatg	gacagcaagc	gaaccggaat	tgccagctgg	7680
15	ggcgccctct	ggtaagggtt	ggaagccctg	caaagttaac	tggatggott	tcttgcgcgc	7740
	aaggatctga	tggcgcaggg	gatcaagatc	atgagcggag	aattaaggga	gtcacgttat	7800
	gacccccgcc	gatgacgcgg	gacaagccgt	tttacgtttg	gaactgacag	aaccgcaacg	7860
	ttgaaggagc	cactcagccg	cgggtttctg	gagtttaast	agctaagcac	atacgtcaga	7920
	aaccattatt	gcgcgttcaa	aagtcgccta	aggtcactat	cagctagcaa	atatttcttg	7980
20	tcaaaaatgc	tccactgacg	ttccataaat	tccctcggt	atccaattag	agtctcatat	8040
	tcactctcaa	tccagatctc	tactctatgc	gagggcccat	gggagcttgg	attgaacaag	8100
	atggattgca	cgcaggttct	cggcgccgtt	gggtggagag	gctattcggc	tatgactggg	8160
	cacaacagac	aatcggtg	tctgatgccg	ccgtgttccg	gctgtcagcg	caggggccc	8220
	cggttctttt	tgtcaagacc	gacctgtccg	gtgcctgaa	tgaactgcag	gacgaggcag	8280
25	cgcggctatc	gtggctggcc	acgacgggcg	ttccttgccg	agctgtgctc	gacgttgtca	8340
	ctgaagcggg	aagggaactg	ctgetattgg	tcgaagtgcc	ggggcaggat	ctcctgtcat	8400
	ctcaccttgc	tccctccgag	aaagtatcca	tactggtga	tgcattgcgg	cggctgcata	8460
	cgttctatcc	ggctacctgc	ccattcgacc	accaagcgaa	acatcgcatc	gagcgagcac	8520
	gtactcggat	ggaagccggt	cttgtcgatc	aggatgatct	ggacgaagag	catcaggggc	8580
30	tcgcgcacgc	cgaactgttc	gccaggctca	aggcgcgcac	gcccgcgcgc	gaggatctcg	8640
	tcgtgaccca	tggcgatgcc	tgettgcgca	atatcatggt	ggaaaatggc	cgcttttctg	8700
	gattcatcga	ctgtggccgg	ctgggtgtgg	cggaccgcta	tcaggacata	gcgttggcta	8760
	cccgtgatat	tgtgaagag	cttggcggcg	aatgttctga	ccgcttcttc	gtgctttacg	8820
	gtatccgcgc	tcccgatctg	cagcgcatcg	ccttctatcg	ccttcttgac	gagttcttct	8880
35	gagcgggacc	caagctagct	tcgacggatc	ccccgatgag	ctaagctagc	tatatcatca	8940
	atztatgtat	tacacataat	atcgactca	gtctttcatc	tacggcaatg	taccagctga	9000
	tataatcagt	tattgaataa	tttctgaatt	taaacttgca	tcaataaatt	tatgtttttg	9060
	cttggactat	aatacctgac	ttgttatctt	atcaataaat	atttaaacra	tatttctttc	9120
	aagatgggaa	ttaattcact	ggccgtcggt	ttacaacgtc	gtgactggga	aaaccctggc	9180
40	gttaccgaa	ttaatcgctt	tgcagcatat	ccccctttog	ccagctggcg	taatagecaa	9240
	gaggcccgca	cggatcgccc	ttcccaacag	ttgcgcagcc	tgaatggcgc	ccgctccttt	9300
	cgttttcttc	ccttctcttc	tgcgccagtt	cgcgcgcttt	ccccgtcaag	ctctaaatcg	9360
	ggggctccct	ttagggttcc	gatttagtgc	tttacggcac	ctcgacccca	aaaaacttga	9420
	tttgggtgat	ggttcacgta	gtgggccatc	gcctgatag	acggtttttc	gccccttgac	9480
45	gttggagtc	acgttcttta	atagtggact	cattgttccaa	actggaacaa	cactcaacc	9540
	tatctcgggc	tattcttttg	atztatagg	gattttgcgc	atctcggaac	caccatcaaa	9600
	caggattttc	gcctgctggg	gcaaaccagc	gtggaccgct	tgtcgcaact	ctctcagggc	9660
	caggcggtga	agggcaatca	gtgtgtgccc	gtctcactgg	tgaagaagaa	aaccaccca	9720
50	gtacattaaa	aaogtccgca	atgtgttatt	aagttgtcta	agcgtcaatt	tgtttacacc	9780
	acaatatatc	ctgccaccag	ccagccaaca	gtcccccgc	cggcagctcg	gcacaaaatc	9840
	accactcgat	acaggcagcc	catcagtcg	ggacggcgtc	agcgggagag	ccgttgtta	9900
	gcggcagact	tctctcatgt	taccgatgct	atccggaaga	acggcaacta	agctgcccgg	9960
	tttgaacac	ggatgatctc	gcggagggtg	goatgttgat	tgtaacgatg	acagagcggt	10020
	gctgctctgt	atcaaatatc	atctccctcg	cagagatccg	aattatcagc	cttcttattc	10080
55	atctctcgt	taaccgtgac	aggctgtcga	tcttgagaac	tatgocgaca	taataggaaa	10140
	tcgctggata	aagccgctga	ggaagctgag	tggcgctatt	tctttagaag	tgaacgttgc	10200
	cgatatcaac	tccccatccc	attgctcacc	gaatgggtaca	ggtcggggac	ccgaagttcc	10260
	gactgtcgcc	tcccgctgat	cccgctgatc	cgaccccaga	tctagatctg	gggctgagaa	10320
	agcccagtaa	ggaacaact	gtaggttcga	gtcgcgagat	cccccggaac	caaagggaagt	10380
60	aggttaaacc	cgctccgato	agggcgagcc	acggccaggcc	gagaacattg	gttctctgtg	10440
	gcacgggat	tggcggtatca	aacactaaag	ctactggaac	gagcagaagt	cctccggccg	10500
	ccagttgcca	ggcggtaaag	gtgagcagag	gcacggggagg	ttgccacttg	cgggtcagca	10560

	cggttccgaa	cgccatggaa	accgcccccg	ccaggccccgc	tgcgacgccg	acaggatcta	10620
	gcgctgcgtt	tggtgtcaac	accaacagcg	ccacgccccg	agttccgcaa	atagccccc	10680
	ggaccgccat	caatcgatc	gggtaccta	gcagagcggc	agagatgaac	acgaccatca	10740
5	gcggctgcac	agcgccctac	gtcgccgcga	ccccgccccg	caggcggtag	accgaaataa	10800
	acaacaagct	ccagaatagc	gaaatattaa	gtgcgcgcag	gatgaagatg	cgcattccacc	10860
	agattccccg	tggaaatctg	cggacgatca	tcacgagcaa	taaaccgccc	ggcaacgccc	10920
	gcagcagcat	accggcgacc	cctcgccctc	gctgttcggg	ctccacgaaa	acgcccggaca	10980
	gatgcgcctt	gtgagcgtec	ttggggccgt	cctcctgttt	gaagaccgac	agcccaatga	11040
10	tctcgccgtc	gatgtaggcg	ccgaatgcca	cggcatctcg	caaccgttca	gcgaacgcct	11100
	ccatgggctt	tttctectcg	tgctcgtaaa	cggacccgaa	catctctgga	gctttcttca	11160
	gggcccacaa	tcggatctcg	cggaaatcct	gcacgtcgge	cgtccaage	cgtcgaatct	11220
	gagccttaat	cacaattgtc	aatttttaate	ctctgtttat	cggcagttcg	tagagcgcg	11280
	cgtgcgctccc	gagcgatact	gagcgaagca	agtgcgtcga	gcagtgcccg	cttgttccctg	11340
15	aatgcccagt	aaagcgctgg	ctgctgaacc	cccagccgga	actgacccca	caaggcccta	11400
	gcgtttgcaa	tgacccaggt	catcattgac	ccaggcggtg	tccaccagge	cgtgcctcg	11460
	caactcttcg	caggcttcgc	cgacctgctc	gcgccacttc	ttcaccgggg	tggaatccga	11520
	tccgcacatg	aggcggaagg	tttccagctt	gagcgggtac	ggctcccggg	gcgagctgaa	11580
	atagtcgaac	atccgtcggg	ccgtcggcga	gagcgttcgg	tactttctcc	atatgaattt	11640
20	cgtgtagtgg	tcgccagcaa	acagcacgac	gatttccctc	tcgatcagga	cctggcaacg	11700
	ggacgttttc	ttgccacggg	ccaggacgcg	gaagcgggtg	agcagcgaca	ccgattccag	11760
	gtgcccacag	cggctcgagc	tgaagcccat	cgccgtcgcc	tgtaggcgcg	acaggcattc	11820
	ctcgcccttc	gtgtaatacc	ggccattgat	cgaccagccc	aggtcctggc	aaagctcgta	11880
	gaacgtgaag	gtgatcggt	cgccgatagg	ggtgcgcttc	gcgtactcca	acacctgctg	11940
25	ccacaccagt	tcgtcatcgt	cggcccgagc	ctcgacgcgc	gtgtaggtga	tcttcacgtc	12000
	cttgtttgac	tggaataatga	ccttgttttg	cagcgctcgc	cgcgggattt	tcttgttgcg	12060
	cgtggtgaac	agggcagagc	gggcccgtgt	gtttggcatc	gctcgcatcg	tgtccggcca	12120
	cggcgcaata	tcgaacaagg	aaagctgcat	ttccttgatc	tgctgcttcg	tgtgtttcag	12180
	caacgcgggc	tgttggcct	cgctgacctg	ttttgccagg	tcctcgccgg	cggtttttcg	12240
30	cttctgttcg	gtcatagtcc	ctcgctgtgc	gatggctcgc	gacttcgcca	aacctgcccgc	12300
	ctcctgttcg	agacgacgcg	aacgctccac	ggcggccgat	ggcgcgggca	gggcaggggg	12360
	agccagttgc	acgctgtcgc	gctcgatctt	ggcgttagct	tgctggacca	tcgagccgac	12420
	ggactggaag	gtttcgccgg	gcgcacgcat	gacgggtcgg	cttgcgatgg	tttcggcatc	12480
	ctcggcgga	aaaccccgct	cgatcagttc	ttgcctgtat	gccttcgggt	caaacgtccg	12540
35	attcattcac	cctccttgcg	ggattgcccc	gactcacgcc	ggggcaatgt	gccccttatc	12600
	ctgatttgac	cgccttggtg	ccttggtgtc	cagataatcc	accttatcgg	caatgaagtc	12660
	ggtcccgtag	accgtctggc	cgctcttctc	gtacttggta	ttccgaatct	tgccctgcac	12720
	gaataccagc	gaccccttgc	ccaaataact	gcctggggcc	tcggcctgag	agccaaaaca	12780
	cttgatgcgg	aagaagtcgg	tgcgctcctg	cttgtcgccg	gcacgttgg	gccacatcta	12840
40	ggtactaaaa	caattcatcc	agtaaaatat	aatattttat	tttctcccaa	tcaggcttga	12900
	tccccagtaa	gtcaaaaat	agctcgacat	actgttcttc	cccgatatcc	tccctgateg	12960
	accggacgca	gaaggcaatg	tcataccact	tgtccgccc	gcccgttctc	ccaagatcaa	13020
	taaagccact	tactttgcca	tctttcacaa	agatgttgc	gtctcccagg	tcgcccgtgg	13080
	aaaagacaag	ttcctottcg	ggcttttccg	tctttaaaaa	atcatacagc	tcgcgaggat	13140
45	ctttaaatgg	agtgtcttct	tcccagtttt	cgcaatccac	atcgccgaga	tcgttattca	13200
	gtaagtaatc	caattcggtc	aagcggctgt	ctaagctatt	cgtataggga	caatccgata	13260
	tgctgatgga	gtgaaagagc	ctgatgcact	ccgcatacag	ctcgataatc	ttttcagggc	13320
	tttgttcate	ttoatactct	tccgagcaaa	ggacgccatc	ggcctcactc	atgagcagat	13380
	tgctccagcc	atcatgccc	tcaaaagtgc	ggacotttgg	aacaggcgac	tttccctcca	13440
50	gccatagcat	catgtccttt	tcccgttcca	catcatagg	ggtcccttta	taccggctgt	13500
	ccgtcatttt	taaatatagg	ttttcatttt	ctcccaccag	cttatatacc	ttagcaggag	13560
	acattctctc	cgtatctttt	acgcagcggt	atttttcgat	cagttttttc	aattccgggtg	13620
	atattctcat	tttagccatt	tattatttcc	ttcctctttt	ctacagtatt	taaagatacc	13680
	caaagaagct	aattataaca	agacgaaatc	caattcactg	ttccttgcac	tctaaaacct	13740
55	ttaaataccag	aaaacagctt	tttcaaagtc	gttttcaaa	ttggcgata	acatagtatc	13800
	gacggagccg	attttgaaac	cacaattatg	ggtgatgctg	ccaacttact	gatttagtgt	13860
	atgatgggtg	ttttgaggtg	ctccagtggt	ttctgtgtct	atcagctgtc	cctcctgttc	13920
	agctactgac	ggggtgggtg	gtaacggcaa	aagcaccgcc	ggacatcagc	gctatctctg	13980
	ctctcactgc	cgtaaaacat	ggcaactgca	gttcaactac	accgcttctc	aacccgggtac	14040
60	gcaccagaaa	atcattgata	tggccatgaa	tggcggtgga	tgccgggcaa	cagcccgcat	14100
	tatggcggtt	ggcctcaaca	cgtttttacg	tcaacttaaa	aactcaggcc	gcagtcggta	14160
	acctcgcgca	tacagccggg	cagtgcgctc	atcgtctgcy	cggaaatgga	cgaacagtg	14220
	ggctatgtcg	gggctaaatc	gcgccagcgc	tggctgtttt	acgcgtatga	cagtctccgg	14280

34

5 aagacgggtg ttgcgcacgt attcggtgaa cgcactatgg cgacgctggg gcgtottatg 14340
agectgctgt caccctttga cgtggtgata tggatgacgg atggctggcc gctgtatgaa 14400
tcccgcctga agggaaagct gcacgtaatc agcaagcgat atacgcagcg aattgagcgg 14460
cataacctga atctgaggea gcacctggca cggctgggac ggaagtgcgt gtcgtttctca 14520
aaatcgggtg agctgcatga caaagtcacg gggcattatc tgaacataaa acactatcaa 14580
taagttggag tcattaceca attatgatag aatttacaag ctataagggtt attgtcctgg 14640
gtttcaagca ttagtccatg caagttttta tgotttgccc atttatataga tatattgata 14700
agcgcgctgc ctatgccttg ccccttgaaa tcottacata cggcgatatc ttctatataa 14760
aagatatatt atcttatcag tattgtcaat atattcaagg caatctgcct cctcatcctc 14820
10 ttcatcctct tcgtcttggt agctttttaa atatggcgct tcatagagta attctgtaa 14880
ggtccaattc tcgttttcat acctcggtat aatcttacct atcacctcaa atgggtcgct 14940
gggtttatcg cacccecgaa cacgagcacg gcacccgcga ccactatgcc aagaatgccc 15000
aaggtaaaaa ttgccggccc cgccatgaag tccgtgaatg ccccgacggc cgaagtgaag 15060
ggcaggccgc caccagggc gccgcccctca ctgcccggca cctggtcgct gaatgtcgat 15120
15 gccagcacct gccgcacgtc aatgcttccg ggcgtcgcgc tcgggctgat cggccatccc 15180
gttactgccc cgatcccggc aatggcaagg actgccagcg ctgccatttt tggggtgagg 15240
ccgttcgcgg ccgagggggc cagcccctgg ggggatggga ggcccgcgtt agcg 15294

20

Figure 1




Figure 2

Variety/line	Mother variety	Dm %	Fructose % of Dm	Glucose % of Dm	Sucrose % of Dm
Producent		27,8	1,8	2,2	3,1
Prevalent		27,1	1,2	1,7	2,7
Dinamo		27,7	0,8	1,7	2,2
Kuras		27,9	1,8	2,2	2,4
AM98-2012	Producent	22,3	3,1	4,0	4,7
AM98-2019	Prevalent	18,4	2,9	4,4	3,8
AM98-2021	Prevalent	17,1	3,1	5,9	3,5
AM99-2002	Dinamo	19,3	2,4	3,8	3,0
AM99-2003	Dinamo	18,8	2,5	5,5	3,2
AM99-2004	Dinamo	11,7	4,0	6,8	2,1
AM00-2040	Kuras	21,0	5,0	6,3	3,5
AM00-2041	Kuras	19,3	5,6	7,1	3,0

Figure 3

3 a) pHS1 for gene-inhibition of StGH1

pHS1




3 b) pHS2 for gene-inhibition of StGH2

pHS2



3 c) pHS3 for over-expression of StGH1

pHS3



3 d) pHS4 for over-expression of StGH2

pHS4




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8





Figure 9



Enhanced amylose production in plants

Abstract

- 5 The invention relates to methods for increasing the amylose content in plants, preferably in potato plants, by expressing a starch biosynthesis enhancing protein. The invention furthermore relates to an expression cassette expressing the polypeptide in potato plants, preferably in the tubers, the transgenic plants expressing the polypeptide and to the use of said transgenic plants for the production of fine chemicals especially other
- 10 than native starches.

PCT/EP2004/002096



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.